

AP1650 C. ✓

57587

INSTITUTIONS
OF ASTRONOMICAL
CALCULATIONS.

PART II.

CONTAINING THE
ASTRONOMY and GEOGRAPHY
OF
TRANSITS,
AND THE
PRINCIPLES of CALCULATION;

Illustrated and applied to the ensuing
TRANSIT of VENUS in 1769;

WITH THE
CONSTRUCTION and Use of a TRANSIT GLOBE
FOR THAT PURPOSE.

EMBELLISHED with THREE large COPPER PLATES,
AND A
MAP of the Great SOUTH SEA.

By B. MARTIN.

L O N D O N;

Printed for and sold by the AUTHOR in FLEET-STREET,
(No. 171) and by all Booksellers in City and Country.

1650

*4-191

WELL

1053 see p.

Gift of Mrs Prof Farnham

TRANSFERRED TO
ASTRONOMICAL OBSERVATORY

AP1650

544
45

P R E F A C E.

IN the first Part of this third Volume of *Institutes* having treated largely on the Doctrine of Solar and Lunar ECLIPSES, with the Rationale and Method of calculating the same from the best TABLES extant (which are given at the End thereof); I judged the Theory and astronomical Calculations of a planetary TRANSIT would very naturally follow that Subject; a Transit being in itself really a solar Eclipse, only in a smaller Degree. Besides, not only the Curiosity but Importance of such a Branch of Astronomy, so essentially necessary to the Perfection of the whole Science, could by no Means be omitted in a course of Institutions of this Kind. Add to this, that the Doctrine of TRANSITS is a recent Topic, never mentioned by former Astronomers, and but very slightly touched upon by the Moderns; the Phænomenon having been never but once publicly observed; and those Observations made in so imperfect a Manner, as to leave us rather under more Uncertainty about the Solar Parallax than we were before; and therefore, an Attempt to elucidate so intricate and obscure a Theory, and to supply the arduous Calculations relative to the next Transit in a most general Manner, I think cannot but be acceptable to every one who is a well Wisher to the Prosperity and Honour of his Country, as these are wholly founded in the Improvement of the Sciences in general, but most immediately in the practical Application of Astronomy, Geography, and Navigation.

I have also endeavoured to facilitate the difficult Ideas relative to the Phænomena of a Transit, by showing how

they naturally arise from, and are solvable by the Properties of the Globe, and particularly the terrestrial one; as also by a new Construction of that Globe, by which all the Phases of a Transit, for any given Minute of Time, are instantly exhibited for all the Inhabitants of the Earth; for which Reason it justly merits the Title of a TRANSIT GLOBE.

Besides three large Copper Plates of Figures and Diagrams, as a thorough Illustration of the general Subject, I have added a Map of the Great South Sea, to exhibit a View of the Island in that Part of the World where the Transit may be observed to the greatest Advantage; the Author of it is Mr. Pingrè, who has given a large and particular Account of the Discovery and Condition of those Islands, to which I refer such Readers as understand French. But those who have a Globe (especially the Transit Globe) can have no Occasion for any other geographical Assistance, to render the practical Part of this sublime Study very easy and entertaining.

A LECTURE upon this Subject, which I propose to give, with Experiments on the above-mentioned Instruments, and other new PIECES of MECHANISM, is all that remains, as I apprehend, to render the Doctrine of Transits as universally intelligible, as it can ever be expected.

C O N T E N T S.

C H A P. I.

*The Astronomical ELEMENTS of a TRANSIT explain'd, with
the RULES for predicting such Events.* page I

C H A P. II.

*The ASTRONOMICAL CONSTRUCTION of a TRANSIT upon
the SURFACE of a Celestial GLOBE. The NATURE of
PARALLAXES explain'd, with the general METHOD of
CALCULATION for the CENTER of the EARTH.* 5

C H A P. III.

*The general METHOD of calculating the PHASES and DURA-
TION of a TRANSIT with regard to any particular PLACE
upon the EARTH'S SURFACE, by Means of the PARAL-
LAXES in LONGITUDE and LATITUDE.* 9

C H A P. IV.

*The foregoing METHOD of CALCULATION exemplified and
illustrated, in a COMPUTATION of the PHÆNOMENA of
the ensuing TRANSIT of June 3, 1769, in respect to the
CENTER of the EARTH.* 14

C H A P.

CHAP. V.

The METHOD of calculating any PHASE of a TRANSIT for a particular PLACE, exemplified in a COMPUTATION of the TIME of apparent INGRESS in the TRANSIT of June 3, 1769, as seen at the CITY of LONDON.

18

CHAP. VI.

The METHOD of computing the Maxima and Minima, or the greatest and least EFFECTS of PARALLAX respecting the TIMES of the PHASES and DURATION of the TRANSIT to OBSERVERS in different Parts of the EARTH'S SURFACE.

21

CHAP. VII.

A CALCULATION for those PLACES where the apparent DURATION of the TRANSIT is a Maximum, or a Minimum.

27

CHAP. VIII.

Of the METHOD of finding the SUN'S PARALLAX by observing the least Distance of VENUS from the SUN. The SUPERIORITY of the METHOD of PARALLAXES; and of the PRECISION to which the SUN'S PARALLAX and DISTANCE of the EARTH and PLANETS from the SUN, may be thereby found,

30

CHAP. IX.

The GEOGRAPHY of a TRANSIT illustrated by a PRAXIS on the Terrestrial GLOBE, exhibiting all the Phænomena of the ensuing TRANSIT of VENUS to the various INHABITANTS of the EARTH.

35

CHAP.

C O N T E N T S.

vii

C H A P. X.

The NATURE of a TRANSIT and its various Phænomena explained, by an Orthographical PROJECTION of the EARTH, SUN, and Planet, upon a Perspective PLANE, and a general COMPUTATION of parallatic TIME from thence.

42

C H A P. XI.

Of the ANALOGY and CORRESPONDENCE between the Orthographical PROJECTION, the METHOD by DIAGRAMS, and the PRAXIS by the GLOBE, in regard to the various PHASES of a TRANSIT.

49

C H A P. XII.

Of the TABLES necessary for computing the TIME in which any given QUANTITY of MOTION is performed; and the MOTION produced in any given TIME; illustrated by EXAMPLES of their Use in the Doctrine of TRANSITS, and other PARTS of ASTRONOMY.

54

C H A P. XIII.

The DESCRIPTION and Use of a TRANSIT GLOBE, by which all the PHASES of the TRANSIT may be instantly exhibited for any given TIME and PLACE, by a new Apparatus for that Purpose

58

CONFIDENTIAL

SECRET

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

INSTITUTIONS
OF
ASTRONOMICAL CALCULATIONS.
PART II.

CHAP. I.

*The Astronomical ELEMENTS of a TRANSIT
explain'd, with the RULES for predicting such
Events.*

4002. **T**HE first Element in the Astronomy of a Transit, is the *Motion of the Node* of the Planet's Orbit; Sir ISAAC NEWTON, and other great Names, have declared positively that the Nodes of the Planet's Orbits are absolutely at rest, and Dr. HADLEY spoke dubiously of it in his Dissertation on the last Transit of Venus; but that Transit compared with the preceding one, observed by the famous HORROX in the Year 1639, put the Motion of Venus's Nodes past all
B Doubt.

2 *The ELEMENTS of a TRANSIT.*

all doubt. And it has been since stated in Dr. HALLEY's Tables at $31''$ per Ann. and by Mr. CASSINI it is made $34''$, who supposes (from a Comparison of a great Number of Observations) that Quantity must be very near the Truth.

4003. The Motion of the Nodes being Retrograde, and at the Rate of $34''$ per Ann. the Place of it will be easily ascertained by proper Tables for any given Moment of Time. Since in the last Transit (1761) the Place of the Ω was $8^{\circ} 14' 40''$, it will in 8 Years have retrograded $4' 32''$, therefore in the next Transit, 1769, the Place of the Node will be $8^{\circ} 14' 35' 20''$.

4004. The Reason why the Venereal Transits succeed each other at such different Intervals of Time, will be best understood by first defining the *Transit-Limits* for this Planet; in order to which let A B be part of the Ecliptic; D E part of Venus's Orbit intersecting the Ecliptic in the Node. Ω ; making the Angle A C D (as viewed from the Earth) $= 8^{\circ} 29' 10''$. Then the Semidiameter of the Sun is at this Node $15' 45''$, and that of Venus (as was found in the last Transit) is $29''$, the Sum of both is $16' 14'' = A D$; hence in the Right-angled Triangle A C D, the Side A C is found $= 110' = 1^{\circ} 50'$, the double of which is $3^{\circ} 40' = A B$, the whole Limit of the Transit. (See Fig. 1 and 2.)

4005. The next Thing to be considered is, that a Transit can only happen at a *Conjunction of the Sun and Venus at her interior Node*; therefore the Time of such a Conjunction, or the Interval between two successive ones, must be determined. Then since the Period of the Planet Venus is 224d. 16h. 42', and that of the Earth 365d. 5h. 49'; it is evident, that if Venus and the Earth set out from any Right Line joining their Centers and that of the Sun, the Time that will lapse before they arrive again to such a Right Line will be equal to a Period of the Planet, and so much more as is equal to the Motion of the Earth in the Interval of two such Conjunctions.

4006.

RULES of predicting such EVENTS. 3

4006. Let the Earth's Motion in that Time be called A, then the Periodical Time of Venus is 5392,75 Hours, and that of the Earth is 8765,75 Hours; and we have this Analogy, viz. As 8765,75h. : 5392,75h. :: $360^\circ + A$: A, and by Division of Proportion, $3373 : 5392,75 :: 360 : A$. Whence we find $A = 575,57^\circ =$ the Motion of the Earth between two Conjunctions; from which subduct 360° , there will remain $215^\circ,57' = 7^s 5^\circ 34'$, which is the Distance in the Ecliptic between every two Conjunctions of the Earth and Venus.

4007. Now 'tis evident, when such a Conjunction happens at either Node, the same will happen again at the same Node when the said Difference $215^\circ,57'$ is so often repeated, or multiplied by such a Number, that the Product shall exceed or fall short of a Multiple of 360 by a Quantity less than the abovementioned Transit Limit. This will be best explained by an Example. The Transit in the Year 1761 was occasioned by a Conjunction of Venus and the Sun near the descending Node, which was then in $\nearrow 14^\circ 40'$, and the Conjunction was in $\nearrow 15^\circ 46'$. If you multiply the Number $215,57$ by 5, the Product is 1077,85, which is less than 1080 ($= 360 \times 3$) by $2,15 = 2^\circ 9'$, which taken from $\nearrow 15^\circ 46'$, gives the Place of the Conjunction at the next Transit in $\nearrow 13^\circ 37'$, which is $47'$ within the Limit B E above the Node \oslash , as the last was within $44'$ of the other Limit A D below it.

4008. Therefore it is evident another Transit must be produced at the same Node at the End of the Interval of 5 Conjunctions, or 8 Years nearly, and therefore must happen again in 1769, June 3. And this was also the Case in the Time of HORROX, when the Transit of Venus in 1631 was follow'd by another 8 Years after in 1639, which was the first ever observed, and that only by himself and his Friend Mr. CRABTREE.

TABLE of future TRANSITS.

4009. In like Manner, if the Number $215^{\circ},57$ be multiplied by such a Number as will give a Product that differs from a Multiple of 360° , by 180° , then will the Conjunction of the Earth and Venus fall within the Transit Limits of the opposite Node. Such a Number of Conjunctions is 81; for $215,57 \times 81 = 17461,17$, which exceeds 17280 ($= 360 \times 48$) by 181; and therefore the 81st Conjunction falls but 1 Degree from the opposite Node.

4010. Now 81 Conjunctions take up the Space of 130 Years (for $5 : 8 :: 81 : 130$ nearly) and consequently the Transits of 1631, and 1639, produce at the Interval of 130 Years, the Transits of 1761, and 1769, at the descending Node ☿.

4011. The next Return of these Transits will be in a less Period of Time, viz. 113 Years; so that the Transit in 1761 will again happen in 1874, and the Transit of 1769 will again return in the Year 1882, both of them at the opposite or *ascending Node*, and after 71 Conjunctions of the Earth and Venus.

4012. I shall here exhibit in one View all the Transits of Venus from the Year 1631 to 2368 inclusive, with the Interval of Years between each two, that the Regularity and Order in which they return, may be the better observed.

Anno Dom.		Node.		Month.		Int. of Years.
1631	—	♊	—	Decemb.		
1639	—	♋	—	Decemb.	—	8
1761	—	☿	—	June	—	122
1769	—	☿	—	June	—	8
1874	—	♋	—	Decemb.	—	105
1882	—	♋	—	Decemb.	—	8
2004	—	☿	—	June	—	122
2012	—	☿	—	June	—	8
2117	—	♊	—	Decemb.	—	105
2125	—	♊	—	Decemb.	—	8
2247	—	☿	—	June	—	122
2255	—	☿	—	June	—	8
2360	—	♋	—	Decemb.	—	105
						2013.

A CALCULATION of a TRANSIT in 2368. 5

4013. That it may further appear how accurately these Periods of 5, 71, and 81 Conjunctions bring about the Transits, even to the same Day of the Month for many Ages together, I shall here subjoin a Computation of that which will happen in 2368, the last in the Table, and distant from the present Time 600 Years.

4014.	Sun's Place.	Apogee.
An. Dom. 1768 —	6 11 42 48 —	3 8 46 55
Years 600 —	11 24 26 32 —	10 6 40
Novemb. 26 —	10 25 14 54 —	53
<hr/>		
Sun's Anomaly —	5 1 24 14 —	3 18 54 28
Add Apog. —	3 18 54 28	
<hr/>		
Sun's Place —	8 20 18 42	
4015.	Venus's Place.	Node.
An. Dom. 1768 —	5 6 36 42 —	2 14 33 1
Years 600 —	3 25 11 12 —	5 10 0
Nov. 26 —	5 18 42 55 —	24
<hr/>		
Place of ♀ —	2 20 30 49 —	2 19 43 25
Place of ♂ —	2 19 43 25 —	8 20 18 42
<hr/>		
	0 0 47 14 —	6 0 35 17

Thus it appears that Venus will be then within 47' of the Node ♂, and therefore far enough within the Transit-limit. Also the Sun (or Earth) is but 35' from the ♂, consequently the *Transit* will then be a very considerable one.

C H A P. II.

The ASTRONOMICAL CONSTRUCTION of a TRANSIT upon the SURFACE of a Celestial GLOBE. The NATURE of PARALLAXES explain'd, with the general METHOD of CALCULATION for the CENTER of the EARTH.

4016. **T**HAT the Use of a *Transit of Venus* may the more fully appear, I shall give the *Astronomical Construction* of the

6 The CONSTRUCTION of a TRANSIT

the next Transit as it may be represented upon the Surface of a *Celestial Globe* by the following Diagram of Circles, for the Beginning or Ingress in the Morning, and viewed on the North Side the Orbit of Venus. Let $H Z O N$ be the general Meridian, $H O$ the Horizon, and Z the Zenith of the Observer; P the Pole of the Equinoctial $\mathcal{A} L$; Q the Pole of the Ecliptic $T S K$; R the Pole of the Orbit of Venus $C S V$; S the Place of the Sun and Venus near the Node; $Z S M$ a Vertical passing thro' the Sun; $P S$ a Meridian thro' the Sun; $Z P S$ the Hour Angle from Noon; $Q S$ a Secondary of the Ecliptic to the Sun; $R S$ a Secondary of the Orbit of Venus; $Q P I$ the Solstitial Colure. (Fig. 3.)

4017. Suppose the Time 9 o'Clock in the Morning, then in the Spherical Triangle $Z S P$ there are known the Co-latitude $Z P$, the Angle $Z P S$, and the Sun's Co-declination $P S$, by which you find the Angle $Z S P$.

4018. Then in the Triangle $Q S P$ there are known all the Sides, viz. $P Q = 20^{\circ} 30'$, $Q S$ a Quadrant, and $P S$ the Co-declination of the Sun; also the Angle $P Q S$ is known, as being the Sun's Distance $S I$ from the first Point of *Cancer* I. From these Data the Angle $P S Q$ is found, and thence the Angle $Q S Z$; also the Angle $R S Q$ is known, being equal to that which the Path of Venus makes with the Ecliptic as seen from the Earth; and therefore the Angle $R S Z$ is known, and of Course the Angle $R S P$.

4019. Therefore all the Parts are known necessary for a general Construction of a Transit; but as the whole Doctrine and Use of the Transit depends upon a proper Resolution of the Planet's *Parallax of Altitude* into the *Parallaxes of Longitude and Latitude*, and learning from thence the particular Quantities of Time to be added to, or subtracted from the Times in which the Phænomena of the Transit happen as viewed from the Center of the Earth, in order to have the Times of the Ingress, Egrefs, Duration,

tion, &c. at the Places of Observation; it will be necessary next to give a Definition of those Parallaxes.

4020. Let ES be a Right Line joining the Centers of the Earth and Sun in the Horizon, and P any Object in that Line as a Planet, &c. AE is the Semidiameter of the Earth. The Planet P will appear to an Eye at the Center of the Earth E upon the Surface of the Sun in F , but if viewed from the Surface of the Earth at A , it will be seen upon the Sun at G , and the Angle $AP E$ is the *Horizontal Parallax* of that Planet. (See Fig. 4.)

4021. Suppose this Planet to be Venus in any other Position V in its Orbit above the Horizon ES , then its Place upon the Sun is at B when seen from the Center E , but viewed from the Surface of the Earth at A , it will appear upon the Sun at C ; and the Angle AVE is the *Parallax of Venus* belonging to the Altitude PV , or Angle VEP ; and this *Parallax* AVE is to the *Horizontal Parallax* $AP E$, as the *Co-sine of the Altitude to the Radius*. (1899.)

4022. The Parallatic Angle at P or V is evidently that by which the Semidiameter AE of the Earth is subtended, or under which it appears at the Planet. Join AB , and the Angle ABE is that under which the Earth's Semidiameter appears from the Sun, and is therefore the *Sun's Parallax*, that great and important Object of our present Enquiries and Researches.

4023. The Difference between the *Parallax of Venus* AVE , or BVC , and the *Solar Parallax* ABE is the Angle BAC , and this is called the *Parallax of Venus from the Sun*. In the Triangle AVB , the Proportion of Distance between the Sun AB , and the Planet AV , being given, and the Sun's Parallax ABE assumed, the Parallax of Venus AVE will be found of Course; because $ABE : AVE :: AV : AB$; and from thence the Angle BAC is also known.

4024. In order to adapt these general Principles to particular Cases and Phases of the Transit, let us suppose the
Globe

8 *The CONSTRUCTION of a TRANSIT*

Globe on which the above Construction is made (Fig. 3.) to be 40 Feet in Diameter, then the Degrees in a great Circle will be nearly four Inches long each, and consequently the Disk of the Sun will be two Inches Diameter, which therefore let be represented by the circular Area $BELFH$; S the Center, in which the Lines PS , QS , RS , meet, from the Pole P of the Equator, Q of the Ecliptic, and R of Venus's Orbit. Let LN be part of the Ecliptic near the Node N , and MN the Path of Venus in which she is descending to the Node over the Face of the Sun in the Chord EB . (Plate I. Fig. 5.)

4025. Draw ES and BS ; then the Right-angled Triangles NSD and $CS D$ are similar, the Angle at D being common to both; and the Angle at $N = CS D$ which is known; also the Side DS is known, being the *geocentric Latitude* of Venus at the Moment of the Conjunction; and from thence CS and CD will be found.

4026. The Chord or Transit-line EB is bisected by the Perpendicular RS , and the Lines SB and CS are known, and therefore the Side $CB (= CE)$; and likewise all the Angles in the Triangles CSB , CSE , will become known.

4027. Again, let SV be the Sum, and SG the *Difference* of the *Semidiameters* of the Sun and Planet; these are both known for the Time of the Transit; therefore in the Triangles CSV , CSG , the Sides CV , CG , will be known with all the Angles in each.

4028. Hence also the Lines GV , AI , will be known both in Motion and Time; and also their Parts AB , BI ; and GE , EV . Thus the Lines DV , DE , DG , and DI , DB , DA , are known; and consequently the Times of the External Contact at V , and the internal Contact at G , will be known at the *Ingress*; also the Time of the Middle of the Transit at C ; and at the internal and external Contacts at the *Egress* at I and A ; and of the whole

Duration of the Transit from V to A, as seen from the Center of the Earth.

CHAP. III.

The general METHOD of calculating the PHASES and DURATION of a TRANSIT with regard to any particular PLACE upon the EARTH'S SURFACE, by Means of the PARALLAXES in LONGITUDE and LATITUDE.

4029 **T**O calculate the Phænomena of the Transit as they appear to an Observer at Z upon the Earth's Surface, it is necessary next to ascertain the Quantity of all the Lines PA, PB, PI, PD, PG, PE, PV; and of the adjacent Angles PVN, PBN, PGN, PDN, &c. which it is easy to do, thus; in the Triangle PDS, the two Sides SD and SP, with the included Angle DSP, are known, which therefore give the Side DP, and the Angle SDP; whose Complement PDQ added to the known Angle SDN, or QDM, will give the Angles PDN, and PDM. Then in the Triangle PDG, the Sides PD, DG, and the Angle PDG, are known, which give the other Side PG, and the Angle at G. Thus in all the other Triangles PDE, PDV, PDI, PDB, PDA, there being two Sides and an Angle given, the remaining Side and Angles will be known.

4030. Let PV, RV, ZV, be drawn from the Poles P R Z, (of the Equator, Orbit, and Horizon) to the Planet V; then in the Triangle ZVP, the Side ZP = Co-Latitude; the Side PV has been already found (4029); and the Angle ZPV is the Time from Noon of the Beginning of the Transit, all which are therefore known; hence we find the Zenith Distance ZV of Venus, and the Angle of Position ZVP.

10 *A CALCULATION of the PHASES,*

4031. Having the Zenith Distance ZV of the Planet, the Parallax belonging to that Altitude is known by the Analogy in (4021) let this Parallax be denoted by VX in the Line ZV continued out, therefore when the Planet is seen from the Center of the Earth at V , it will by this Parallax be depressed to X , as viewed by an Observer at Z . Thro' the Point X , and parallel to the *true Path* (or *Transit-Line*) MN , draw the dotted Line mn , which will be the *apparent Path* of Venus, or that in which she appears to move, as seen from the Earth's Surface at Z . (See Plate II. Fig. 1.)

4032. This Parallax VX is the Measure of the Angle VAB in Plate I. Fig. 4. and consequently if the true Quantity thereof could be estimated by any Astronomical Instrument, or Methods hitherto used for that Purpose, the Parallax of the Sun ABE would be found very near the Truth; for in the Triangle AVB , we have $AB : VB :: VAB : ABV$, or ABE .

4033. But no Means hitherto used to determine the Parallax of any Planet have proved successful; and Astronomy can furnish but one other Method more, by which the *Solar Parallax* can be attempted with any Probability of approaching to it near the Truth; and this is by a *Transit of Venus*; for the Transit of *Mercury* does not afford a *Parallax of that Planet from the Sun* sufficient to answer any such Purpose.

4034. In this Transit, the *Parallax of Altitude* VX is estimated by the Effect which it produces in the *different Times of the Phases*; for this Parallax when properly resolved, will shew a Difference in the Time when any *Ingress*, *Egress*, &c. will happen to an Eye placed in the Center of the Earth, and any where upon its Surface at Z , both in regard to the *Latitude* and *Longitude* of that Place.

4035. Thus (by the Principles of *Mechanics*) the Parallax VX is resolvable into the two others, $X\psi$, and $V\psi$; the

the first is a *Parallax in Longitude*, because to a Person at Z the Planet will appear at X, which is distant from the Secondary R V, to the *East* or *West*, by the Quantity of the Line X v; and therefore the Moments of Time in which the Planet will be seen in the two Secondaries passing through V and X, will be different according to the *Longitude* of the Spectator.

4036. The other Part V v is the *Parallax of Latitude*; for while the Planet is seen at V from the Earth's Center, it is depressed in the Secondary R v to the Point v in the apparent Path. But it is evident, if the Planet be in Contact at V, it cannot be so in any other Point v in the same Circle; but the Moment of Contact will be sooner or later than at V, by the Difference of Time the Planet takes to pass from v in the Secondary to the Point Y in the Circle of Contact Y D V, joining the Centers when in Contact at V and Y in the *true and apparent Paths*.

4037. The Times which the Planet takes to pass thro' the Spaces X v and v Y, will be known from a Computation of the Quantity of the said Lines, which is thus. The Angle Z V P is known (4030) also the Angle R V P, = $90^\circ - P V M$, therefore also, the Angle Z V R = X V v; hence in the Right-angled Triangle v V X, having the Hypothenuse V X, and the Angle at V, the Sides X v, and V v become known.

4038. Then for computing the Value of v Y, it is evident, by drawing S Y, we have $Y v = c Y - c v = c Y - C V$. But C V is known, being half the Motion of

the Central Transit; and $c Y$ is $= \sqrt{S Y^2 - S c^2}$ a known Quantity; for $S c = S C - C c = S C - V v$, both which are known. Also S Y is known being the Sum (or Difference) of the Semediameters of the Sun and Venus; and hence v Y is known.

4039. The Motions v X and v Y being known, the Times in which they are performed are known, from the known geocentric Motion of Venus at the Time of the

12 A CALCULATION of the PHASES.

Transit. In all that has been hitherto said, the *Longitude* of the Observer, or of the Point Z, is its Distance *East* or *West*, from the Secondary R V; and the *Latitude* thereof is its Distance, *North* or *South*, from the Orbit of Venus.

4040. Therefore the *Motion in Longitude* arising from the *Parallax of Longitude* is to be reckoned from or towards the Secondary R V, in the true Path (or Orbit) M N; but the Motion in Latitude, resulting from the *Parallax of Latitude*, is considered in the apparent Path *mn*. We shall now exhibit, in four Diagrams, the *eight* different Phases of the INGRESS and EGRESS of the Planet upon the Disk of the Sun in the next Transit, which will be found, if duly considered, to comprehend all the Varieties incident to that wonderful Phænomenon; the Construction being the same for the interior Contact at every *Ingress* or *Egress*, requires not to be drawn, as it would too much embarrass the Diagrams with unnecessary Lines.

4041. When the INGRESS is viewed by an Observer at Z, on the *North Side* of the Planet's Orbit, in the *Eastern Hemisphere*, or *Forenoon*, then the Construction of it will be such as we have described (in Fig. 1. Plate II.) Wherein it is evident, that ψ X is the Motion in Longitude, and ψ Y, that in Latitude; and the Times of passing these Lines or Spaces, are those by which the true Time of the Ingress at V will be varied, by the Addition or Subtraction of their Sum or Difference, to or from the said Time of Ingress viewed from the Center of the Earth: Whence will arise the following Cases.

4042. *First*; When the Observer is situated upon the Orbit of Venus (or rather the Parallel she describes) he has then no Latitude; the Point ψ coincides with V; and the *Parallax of Altitude* is wholly a *Parallax of Longitude* in the true Path M N.

4043. *Secondly*; If the Observer's Situation be in the Planet's prime Vertical R V, he has then no Longitude, and of Course there will be no *Parallax of Longitude*, but
of

and DURATION of a TRANSIT. 13

of *Latitude* only, viz. Vv ; the Line or Azimuth ZX , coinciding with Rv , makes $vX = 0$; and vY is the only Effect of Parallax in this Case.

4044. *Thirdly*; A Situation any where between the Orbit and its Secondary RV , will give a Parallax of Altitude VX in the Azimuth ZV , productive of a Parallax of *Eastern Longitude* vX , and a Parallax of Latitude Vv .

4045. *Fourthly*; When the Spectator is between the Secondary RV , and the Circle of Declination PV , the Parallax of Longitude will be on the *West Side* of the Circle RV . These are all the Cases which require to be specified for the Ingress, at present, in regard to the Position of the Observer.

4046. The next Thing is to consider how the *Time of Ingress* is affected by these different and variable Parallaxes. The true Time of Ingress is in the Secondary RV , when the Planet is at V ; the Time of apparent Ingress is the Moment of the Planet's being at Y in the apparent Path. Hence also will arise four Cases in regard to the Time.

4047. *First*; When the Point X falls *Eastward* of the Circle RV , it is evident the Time of the Ingress will be *retarded* by all the Time required in passing through the Space Xv , or Parallax of Longitude.

4048. *Secondly*; When the Point X falls to the West of the Circle RV , the Time of Ingress will be accelerated by the Parallax of Longitude.

4049. *Thirdly*; When the Point Y (in which the Center of the Planet is at the Moment of apparent Ingress) falls *Eastward* of RV , the Time of Ingress will be accelerated by the Parallax of Latitude vY .

4050. *Fourthly*; But when the Point Y is on the *West Side* of the Circle RV , then the Time of Ingress will be retarded by the Time the Planet takes in passing from v to Y .

All these Cases are general, and hold good for all Ingresses,

gresses, Egresses, interior Contacts, &c. wheresoever observed.

4051. From what has been premised, it easily appears (1) That when X and Y fall on different Sides of R *v* the Time of Ingress, Egress, &c. will be *accelerated* or *retarded* by the Sum of both the Parallaxes, or by the Time of passing over $Xv + vY = XY$. (2.) That when the Points X, Y, fall both on the same Side the Circle R *v*, then the *Time of Ingress*, &c. will be *accelerated* or *retarded* by the Difference of the Parallaxes, or by the Time of passing over $Xv - Yv = XY$.

4052. And since only the same Lines ZX, R *v*, and P V; and the same Points X, *v*, Y, are concerned in the Ingress, Egress, &c. of the Planet, as viewed in the *Eastern* or *Western Hemisphere*, or on the *North* or *South Side of the Orbit*; therefore it will be evident, by Inspection only, when the Time resulting from the Parallaxes is *to be added to*, or *subtracted from the true Time*, in order to have the *Moment of visible Ingress*, or *Egress*, or *Quantity of apparent Duration*. And to make every Thing in this intricate Doctrine clear and easy, we shall next illustrate and apply the foregoing Theory of a Transit, by a Calculation of that which is next to happen, for the Time of its *Ingress*, as seen from the Center of the Earth.

CHAP. IV.

The foregoing METHOD of CALCULATION exemplified and illustrated, in a COMPUTATION of the PHÆNOMENA of the ensuing TRANSIT of June 3, 1769, in respect to the CENTER of the EARTH.

4053. **I**N Order to this, the Places of the Sun and Venus must be found for some particular Point of Time, near the Middle of the Transit, but it may be either before or after.

We

of the ensuing TRANSIT of VENUS. 15

We shall here chuse the Moment of *Eight o'Clock*. From MAYER's Solar Tables (in the preceding Part of these *Insituts*) the Place of the Sun is had as below.

	☉ Long.	Apogee.
An. 1769	9 10 24 12	3 8 56 52
June 3	5 1 47 23	27
Hour 8	19 43	
	<hr/>	<hr/>
	2 12 31 18	3 8 57 19
Equat. add	50 7	2 12 31 18
	<hr/>	<hr/>
The Sun's true Place	2 13 21 25	11 3 33 59

4054. From Dr. HALLEY's Tables of Venus's Motions, her Place for the same Time is had as follows.

	Long of ♀	Aphel.	Node ♄
1769	0 23 0 19	10 7 36 26	2 14 33 32
May 23	7 19 6 36	17	10
Hour 8	32 3		
	<hr/>	<hr/>	<hr/>
	8 12 38 58	10 7 37 13	2 14 33 42
Equat. add	39 6	8 12 38 58	6
	<hr/>	<hr/>	<hr/>
	8 13 18 4	10 5 1 45	8 14 33 42
Reduct. add	7	Mean Anomaly.	Place of ♄
	<hr/>		
Place in the Ecliptic.	8 13 18 11		

4055. The Place of the Earth is - - - 8 13 21 25
From which subduct the Place of ♀ 8 13 18 11

There remains the Difference - - 3 14

4056. The hourly Motion of Venus in the Ecliptic will (being so near the Node) be the same with that in her Orbit, viz. 4'. The hourly Motion of the Earth at the Time of the Transit is 2' 23", 5 as will appear from any Ephemeris.

If then we say, as the Difference of the hourly Motions $1' 36''.5 = 96''.5$ is to the Difference of their Places $3' 14'' = 194''$ so is the hourly Motion of Venus $4' = 240''$ to her Distance from the Plane of Conjunction $482''.5 = 8' 2''.5$. This being added to Venus's Place, gives the Point $8^s 13' 26' 13'' \frac{1}{2}$ wherein the Conjunction happens.

4057. The Time in which Venus passes over the Space of $482''.5$ is thus found: As $240'' : 60' :: 482''.5 : 120'.63 = 2^o 0' 38''$; this added to the given Time of 8 o'Clock, gives $10^o 0' 38''$ for the mean Time of Conjunction, at the Center of the Earth.

4058. From the Place of the Node $\vartheta = 8 14 33 42$
 Subtract the Place of Conjunction $8 13 26 13.5$

There remains the Distance from $\vartheta = 1 7 28.5$

4059. The Inclination of Venus's Orbit to the Plane of the Ecliptic, being $3^o 23' 20''$, her heliocentric Latitude at the Conjunction will be $= 4'$.

4060. The mean Distance of the Earth from the Sun being put $= 10000$, it's Distance at the Time of the Transit will be 10151 ; the Distance of Venus from the Sun will be 7262.6 ; and her Distance from the Earth 2888.5 . Therefore say: As $2888.5 :: 7262.6 :: 4' : 10' 3'' =$ Venus's geocentric Latitude at the Conjunction; and is the Measure of the Line S D in the Diagrams of the Transit.

4061. The Inclination of Venus's Orbit to the Ecliptic will likewise be magnified in a *geocentric View*, in the inverse Ratio of the Distances of the Sun and Earth; therefore say: As $2888.5 : 7262.6 :: \text{Tangent of } 3^o 23' 20'' : \text{Tangent of } 8^o 28' 6'' = \text{M N L}$, which is also the Quantity of the Angle R S Q. (Fig. 5. Plate I.)

4062. The Obliquity of the Ecliptic being $23^o 29'$, and the Sun's Place therein $73^o 26' 13''$ found as above (4053.) You from thence find the Sun's Declination $23^o 27' 15'' \frac{1}{2}$; and its Co-declination is therefore S P $= 67^o 32' 44'' \frac{1}{2}$.

of the TRANSIT for the EARTH'S CENTER. 17

4063. In the Spherical Triangle PSQ all the Sides are known; and from thence you find the Angle $PSQ = 7^{\circ} 11' 37'' \frac{1}{2}$; This added to the Angle RSQ , gives the Angle $RSP = 15^{\circ} 39' 43'' \frac{1}{2}$.

4064. In the Triangle SDC , there are known the Side $SD = 10' 3'' \frac{1}{2}$, the Angle $DS C = 7^{\circ} 11' 37'' \frac{1}{2}$, and the Right Angle at C ; from which you find the Side $CD = 1' 29''$, and the Side $CS = 9' 57''$. Also the Angle $CDS = 81^{\circ} 31' 54'' = QDM$.

4065. In the Right-angled Triangle SCV , there are known the two Sides $SC = 9' 57''$, and $SV = 16' 14'' = 15' 45'' + 29''$, Sum of the Semidiameters of the Sun and Venus; and from thence we have the Side $CV = 12' 49''$; and therefore $CV - CD = DV = 11' 20'' \frac{1}{2}$. Also the Angle $CVS = 37^{\circ} 47' 46''$, and the Angle $CSV = 52^{\circ} 12' 14''$.

4066. In the Triangle CSG we have given the Side CS , and the Side $SG = (15' 45'' - 29'' =) 15' 16'' =$ the Difference of the Semidiameters of the Sun and Venus; from which we find the Side $CG = 11' 34'' \frac{1}{2}$; whence also $DG = 10' 5'' \frac{1}{2}$.

4067. The heliocentric Motion of Venus from the Earth is (as before observed) $1' 36'' \frac{1}{2} = 96'',5$; and her geocentric Motion from the Sun will exceed that in the Ratio of 7262,6 to 2888,5; hence her apparent Motion upon the Sun's Disk will be $4' 2'' \frac{1}{2}$ per Hour.

4068. Therefore $242'' \frac{1}{2} : 60' :: 769'' \frac{1}{2} : 3^h 10' 19''$, the Time of Venus passing through CV , which therefore is the *Half-duration of the Transit*, and so the *Whole Duration* would be $6^h 20' 38''$, if seen from the Center of the Earth.

4069. The Time of her passing through DV will in like Manner be found $= 2^h 48' 19''$, which taken from the Time of the Conjunction $10^h 0' 38''$, leaves $7^h 12' 19''$ for the *Time of the Beginning, or external Contact* at V .

4070. The Time of passing through GD is found by
D
the

18 *A CALCULATION of the TRANSIT*

the same Analogy to be $2^h 29' 43''$, which subducted from the Time of passing through D V will leave $18' 36''$ for the Time between the *external Contact* at V, and the *internal Contact* at G, which therefore will happen at $7^h 30' 55''$.

C H A P. V.

The METHOD of calculating any PHASE of a TRANSIT for a particular PLACE, exemplified in a COMPUTATION of the TIME of apparent INGRESS in the TRANSIT of June 3, 1769, as seen at the CITY of LONDON.

4071. **W**E next proceed to the Calculation of the Phenomena of the Transit for any particular Place upon the Surface of the Earth, thereby to find what Differences will arise from the Effects of *Refraction* in regard to the Times of their Appearance and Duration. This I shall exemplify in a Computation of the BEGINNING of the Transit, for the exterior and interior Contact, as they will be observed by a Spectator at the City of LONDON, denoted by Z. (Fig. 5. Plate I.)

4072. Then in the Triangle D P S there are known the two Sides S P and S D, and the included Angle at S; by which we find the other two Angles $S D P = 172^\circ 47' 52'' \frac{1}{2}$ and $D P S = 1' 22'' \frac{1}{2}$; and also the Side D P $= 67^\circ 23' 8''$. The Complement of S D P to 180° is the Angle P D Q $= 7^\circ 12' 7'' \frac{1}{2}$, which being taken from Q D M ($= C D S = 81^\circ 31' 54'' \frac{1}{2}$) there will remain the Angle P D M $= 74^\circ 19' 47''$.

4073. Hence in the Triangle P D G, there are given the two Sides D P, D G, and the included Angle D; by which we find the Side P G $= 67^\circ 20' 25''$. Also in the Triangle P D V, the Sides P D and D V are known, and the same Angle at D; therefore we find the Side P V $= 67^\circ 20' 4''$. Again, in the same Triangles, are found the

the external Angles $PGM = 74^{\circ} 23' 5''$, and $PVM = 74^{\circ} 23' 34''$.

4074. In the Triangle ZPV are known the Side $ZP = 38^{\circ} 29'$; the Side $PV = 67^{\circ} 20' 4''$; and the included Angle $ZPV = 108^{\circ} 4' 45'' = 7^h 12' 19''$ the Time from Noon (4069) of the beginning of the Transit. From thence we find the Side $ZV = 83^{\circ} 12' 48''$, the Distance of Venus from the Zenith of LONDON. Her *Altitude* therefore at that Moment is $6^{\circ} 47' 12''$.

4075. In like Manner the Moment of interior Contact, being $7^h 30' 55''$, gives the Angle $ZPG = 112^{\circ} 43' 45''$, &c. then with the Side ZP and PG , we find the third Side $ZG = 85^{\circ} 25' 42''$, the Zenith Distance of the Planet at G .

4076. In the same Triangles we find the Angle $ZVP = 36^{\circ} 4'$, and the Angle $ZGP = 35^{\circ} 9'$. If the Angle PVM be taken from the Right Angle $RV M$, there will remain the Angle $RV P = 15^{\circ} 36' 26''$, which added to the Angle ZVP , gives the Angle $RV Z = 51^{\circ} 40' 26''$. In the same Manner you find the Angle $RGZ = 50^{\circ} 45' 55''$. (See Fig. 2. Plate II.)

4077. If we now assume the *Sun's Parallax* $AGE = 10''$, we shall have the horizontal Parallax of Venus $AP E = 35'',143$; and from thence the Difference $= 25'',143 = FE G$, the horizontal Parallax of Venus from the Sun. (See Fig. 4. Plate I.)

4078. Hence the Parallax belonging to the Zenith-distance ZV will be found $= 24'',966$; and that for the Distance $ZG = 25'',063$; therefore $25''$, which is the Mean, may be used for either of them indifferently without Error, and so we have $VX = 25''$. (See the Diagrams, Plate II.)

4079. Therefore in the Right-angled Triangle vVX , there are known the Side VX , and the Angle $XV v = RVZ$; by these are found the Side $Xv = 19'',61$; which is the *Parallax in Longitude*; and the Side Vv

20 *A CALCULATION of the PHASES*

= 15",503, which is the Parallax in Latitude. And because $Cc = Vv$, therefore $Sc = 9' 41'',57$. (4064.)

Therefore $\sqrt{SY^2 - Sc^2} = cY = 13' 1'',32$ from which take $CV = 12' 49'',5$, and there will remain $vY = 11'',82$ (4038.)

4080. This Method of finding the Value of vY is indeed strictly mathematical; but the following is much easier, and very exact. The Arch VY may be taken for Part of the Tangent to the Circle DVL in the Point V ; also the Angles $vVX + vVY = 90^\circ = vVX + SVC$; therefore the Angle $vVY = SVC$; and is given by (4065.) Hence the Parallax of Latitude Vv being known, its Effect vY is known also by this Analogy. As the Radius to the Tangent of vVY , so is Vv to vY .

4081. The hourly Motion of Venus from the Sun is $242'',5$ as we have shewn (4068) therefore she will pass over the Line or Space $vX = 19'',61$ in $4' 51'',12$ of Time, which will be the Effect of the *Parallax in Longitude*. Also the Time in passing over the Space Yv will be $2' 55'' 44$; and this is the Effect of *Parallax in Latitude*. Therefore the whole Effect of Parallax is the whole Time of passing through YX , which is $7' 46'' \frac{1}{2}$; and by so much will the Time of the Ingress or external Contact happen sooner to an Observer at *London Z*, than to an Eye placed in the Earth's Center; and therefore we may expect to see the Transit begin at $7^h 4' 32'', \frac{1}{2}$ in the *Evening*.

4082. In the same Manner it will be found, that the Acceleration of the Time of the *internal Contact* at G will be $4' 47''$ by the Parallax of Longitude, and $3' 20''$ by that of Latitude. So the whole Acceleration will be $8' 7''$, and therefore we may expect the interior Contact to happen at $7^h 22' 48''$. Hence the Time between the external and internal *apparent Contacts* will be $18' 16''$; shorter than the same Interval viewed from the Earth's Center by $20''$, (4068.)

C H A P.

CHAP. VI.

The METHOD of computing the Maxima and Minima, or the greatest and least EFFECTS of PARALLAX respecting the TIMES of the PHASES and DURATION of the TRANSIT to OBSERVERS in different Parts of the EARTH'S SURFACE.

4083. **W**E have shewn that there will be considerable Differences between the *true and apparent Times* of the Phases and Durations of the Transit, and how they may be estimated at any particular Place; but we are further to remark, that these Differences are *greater or less*, according to the different Circumstances or Situation of the Observer with regard to *Latitude and Longitude*; the *Eastern or Western Hemisphere*, the *North or South Side of the Orbit*, &c. And it will appear that there is one Situation of the Observer which will make the Difference of Time between his observed Phase and the central one the greatest possible, or a *Maximum*; and consequently these Places are the principal Objects of our present Enquiry.

4084. This *Maximum*, 'tis evident, consists in the largest Quantity of the Line YX , which again depends upon a particular Quantity of the Angle XVv , or rather of the Angle XVY ; for since the Angle VYX , and the Line XV are given Quantities, the Case is reduced to this Problem, viz. to adapt a given Line XV in a given Angle Y , so as to make the Side YX a *Maximum*. To solve which, let the two equal Lines XV and AB have two different Positions in the given Angle VYX (Fig. 6. Plate I.) then, by Trigonometry, we have $AB : \text{Sine of } Y :: AY : \text{Sine of } B$; and $XV : \text{Sine of } Y :: XY : \text{Sine of } V$; therefore we have $AY : B :: XY : V$; from whence it appears that the Side AY encreases with the Sine of the Angle B , and is a *Maximum* when that Sine is greatest; that is, when

22 A CALCULATION of the PHASES

when the Point B arrives at V, where the Angle X V Y is a right one, and the Side X V, of Course, a *Maximum*.

4085. Hence then, (in the Diagrams of Plate II.) it is evident, the Secondary Z V, and the Radius S V, must, in this Case, coincide, and make one right Line; or in other words, the *Secondary passes thro' the Centers of Venus and the Sun, when the whole Effect of Refraction is the greatest possible*; therefore the Angle Z V R = v V X = V S C = $52^{\circ} 12' 14''$, at the ensuing Transit.

4086. Again, the greater the Zenith Distance Z V is, the greater will be the Refraction V X in Altitude, and therefore Observations of any particular Phase made as near the Horizon as conveniently can be, will be so much the better; we shall therefore suppose that the least Altitude that will admit of a distinct View of the Contact of the Limbs of Venus and the Sun, is 7 Degrees; and in that Case the Refraction X V = $25''$, extremely near. (4021.)

4087. In the next Place, we are to determine from the above Principle, where those two Places are, in one of which the Time of any Phase is *accelerated* as much as possible, and in the other, it is in the same Manner *retarded*; for in these Places there will be the greatest Difference of Time in the Observations that can be obtained.

4088. By the Construction of the Transit (Plate II. Fig. 2.) it appears that the Ingress will be *accelerated* by the Sum of both Parallaxes Y X in the *Afternoon* to an Observer on the *North Side of Venus's Orbit*. And in (Fig. 3.) it is evident, the same Phase will be *retarded* equally from the same Cause to an Observer on the *South Side* of the said Orbit in the *Morning*. The Latitude and Longitude of each of these Places, is therefore to be determined by the following Process.

4089. In the first Case, because the Angle R V Z = $52^{\circ} 12' 14''$ for a *Maximum*; therefore its Complement Z V M = $37^{\circ} 47' 46''$ subducted from the Angle P V M = $74^{\circ} 23' 34''$ will leave the Angle P V Z = $36^{\circ} 35' 48''$.

Again,

when of the greatest and least QUANTITY. 23

Again, by Supposition, the Side $ZV = 83^\circ$; and the Side $PV = 67^\circ 20' 4''$. Therefore in the Triangle PVZ , having given two Sides and the included Angle, we find the third Side $ZP = 38^\circ 31' 43''$ whose Complement is the *Latitude of the Place Z* $= 51^\circ 28' 17''$, which is the first Thing required.

4090. Then for the *Longitude of the Place Z*, we find in the same Triangle, the Angle $ZPV = 108^\circ 10' 57''$, which is the *Longitude of the Place Z*, with regard to the Meridian PV , which passes thro' Venus at the Beginning of the Transit. But since that Angle with respect to *London*, is $108^\circ 4' 45''$; it is plain, the *Meridian of the Place Z* is but $6' 12''$ eastward of the *Meridian of London*. And therefore since the Latitude and Longitude of the Place Z , is so very little different from the Latitude and Longitude of *LONDON*; *this great City has happily the very best Situation that possibly can be to observe the most favourable Ingress that can, in any Transit, ever happen.*

4091. It remains now to find the Place where the same Phase or Ingress is *retarded as much as possible by parallatic Time*. And this (as is evident) from (Plate II. Fig. 3.) will be found on the *South Side of the Orbit*, and in the *Eastern Hemisphere*, or *Morning Observation*.

4092. To compute the Quantity of this *maximum Retardation*, you proceed as follows; the Angle $vVX = rVZ = 52^\circ 12' 14''$ as before; also the Angle $pVN (= PVM) = 74^\circ 23' 34''$, from which if you deduct $ZVN = 37^\circ 47' 46''$, there will remain the Angle $pVZ = 36^\circ 35' 48''$, as in the *London Ingress*.

4093. Then in the Triangle pVZ , the Side $pV = 112^\circ 39' 56''$ (being the Complement of PV in the other Case;) also the Side $ZV = 83^\circ$, by Supposition. Therefore, by having these two Sides, and the included Angle at V , you find the third Side $pZ = 46^\circ 29' 27''$, whose Complement $43^\circ 30' 33''$ is the *South Latitude* of the Place Z of Observation.

24 A CALCULATION of the PHASES

4094. The *Longitude of the said Place* is known from the same Data; for thereby the Angle $V p Z$, will be found $= 54^{\circ} 44' 30''$, which shews the Position of the Meridian passing through Z is so much to the *West* of that passing through the Planet V at the Moment of Ingress; but this through Venus is distant *Westward* from London $108^{\circ} 4' 45''$, as we have shewn (4075.) Therefore, the *Longitude sought* of the Place Z from London, is $162^{\circ} 49' 15''$ *Westward*.

4095. The Quantity of the Angle $Z p V = 54^{\circ} 44' 30''$ turned into Time, is $3^h 38' 58''$, which is the Time from Noon, and therefore subtracted from XII, leaves $8^h 21' 2''$, for the Time of the Morning when the Central Ingress happens at that Place, viz. upon the 4th Day of June, at $21' 2''$ after 8 o'Clock.

4096. Then for the Difference between the central and apparent Time, it will be nearly the same as at London, for in the Triangle $X V v$, the Side $V X$ and the Angle at V are the same here as they were there; and consequently the Sides $X v$, and $V v$, will be found the same, viz. $X v = 19'',61$; and $V v = 15'',5$; but because $S c$ is here $= S C + V v$, we shall find $c Y = 12' 37''$, which taken from $C V = 12' 49'' \frac{1}{2}$, leaves $Y v = 12'' 2$, not $\frac{2}{3}$ of a Second larger than before at the London Ingress. Therefore the Ingress will be retarded here by $7' 46'' \frac{1}{2}$, and so the Time of observing it at the Place Z , will be $8^h 28' 48'' \frac{1}{2}$ in the Forenoon.

4097. The Process is the same in calculating the Times of a *retarded and accelerated Egress*; for it appears that the Egress will be *retarded* by the joint Effect of both Parallaxes, when viewed in the Morning on the North Side of the Orbit of Venus; and the same Egress will be in the same Manner *accelerated* when observed in the Afternoon on the South Side of the Orbit, as in Fig. 1. and 4. Plate II.

4098. To this End the same Lines and Angles must be calculated for the Place of Venus at A , in her external Contact with the Sun's Limb at her *going off*, as was done

at V for her *coming on*, and these will be had from the same Data. Thus we find $AP = 67^{\circ} 27' 2''$; the Angle $PAM = 74^{\circ} 13' 15''$; its Supplement $PAN = 105^{\circ} 46' 45''$, $PAR = 15^{\circ} 46' 45''$. The *Maximum* Angle $ZAR = \angle VAX = 52^{\circ} 12' 14''$, and therefore the Angle $ZAP = 67^{\circ} 58' 59''$; also the Side $ZA = 83^{\circ}$, as before.

4099. Then in the Triangle ZAP , there are given the two Sides AP and AZ , with the included Angle ZAP , by which we find the the third Side $ZP = 67^{\circ} 1' 23''$, whose Complement is $22^{\circ} 58' 37''$, the *Latitude North*, for the Place of the Observer at Z .

4100. In the same Triangle we find the Angle $APZ = 88^{\circ} 11' 30''$, which is the Longitude of the Meridian of the Observer *Westward* from that which passes through the Planet at A . This converted into Time, is $5^h 54' 46''$, which taken from XII, leaves $6^h 5' 14''$, for the Time in the Morning of *June* the 4th, when the *Transit ends there*, as seen from the Center of the Earth.

4101. And because the Angle $\angle VAX$, and the Line AX , are here of the same Value as at the Ingress V , the parallatic Time will be the same, viz. $7' 46'' \frac{1}{2}$; therefore the observed Time of the Egress, or exterior Contact, will be *13' after Six* in the Morning.

4102. In order to find the Longitude of the Place Z from the Meridian of *London*, it must be remembered, that at the Beginning of the Transit in V , the Distance of the Meridian of *London* was $108^{\circ} 4' 45''$ from that of Venus *Eastward*; and that while Venus moves from V to A , which is in Time $6^h 20' 38''$ (4068.) the Meridian of *London* will have passed through $95^{\circ} 9' 30''$ of the celestial Equator, still more eastward; the Distance thereof from that of Venus at the End of the Transit at A will be $203^{\circ} 14' 15''$ *Eastward*, or $156^{\circ} 45' 45''$ *Westward*. But the Longitude of Z is $88^{\circ} 11' 30''$ *Westward* of that of

E

Venus

26 A CALCULATION of the DURATION

Venus also; therefore it is Eastward of London by $68^{\circ} 34' 15''$.

4103. Again (by Fig. 4) it appears that both the Parallaxes conspire to *accelerate the Egress* to an Observer on the *South Side* of Venus's Orbit, viewing it in the *Afternoon*. To determine the *Latitude* and *Longitude* of his Situation for making the Observation to the greatest Advantage, we have $r A Z = 52^{\circ} 12' 14''$, and $A Z = 83$, as before; also $p A M = 105^{\circ} 46' 45''$, therefore $p A r = 15^{\circ} 46' 45''$; consequently $Z A p = 67^{\circ} 58' 59''$. Also $A p = 112^{\circ} 32' 58''$. Therefore in the Triangle $Z A p$, we have known the two Sides $A p$ and $A Z$, with the included Angle at A, by which is found the third Side $Z p = 72^{\circ} 43' 27''$, whose Complement $17^{\circ} 16' 33''$ is the *South Latitude* sought.

4104. By the same Triangle and Data, we find the Angle $A p Z = 74^{\circ} 30'$, which is the Distance of the Meridian of the Observer from that of Venus, to the East. But the Meridian of Venus is then $156^{\circ} 45' 45''$ eastward of London (4102) therefore the Meridian of the Observer at Z is $231^{\circ} 15' 45''$ eastward from LONDON, or $128^{\circ} 44' 15''$ to the *West* of it.

4105. The same Angle $A p Z$ turned into Time, gives $4^h 58'$ from Noon, for the Time of the *central Egress* at that Place; and since all Things are the same here in the parallactic Triangle $v A X$, as in the foregoing Cases, the Time of the Egress will be accelerated by the same Quantity, viz, $7' 46'' \frac{1}{2}$, and therefore the Moment of the observed Contact at Y will be $4^h 50' 14'' \frac{1}{2}$.

4106. As to a Calculation of the Latitudes and Longitudes of Places for the *interior Contacts* at G and B, it is no Ways necessary, for the Latitude will be nearly the same for both Contacts at the *Ingress* and *Egress*; and the Longitude for the interior Contact will be but $4^{\circ} 30''$ different from that of the external one, as it answers to the Time $18' 36''$ of passing through V G and B A; (Fig. 5. Plate II.)
there=

when a Maximum and a Minimum. 27

therefore if $4^{\circ} 30''$ be added to the Longitudes already found for the exterior Contacts at the Ingress, and subducted from those at the Egress, the Longitudes for the interior Contacts will be had.

CHAP. VII.

A CALCULATION for those PLACES where the apparent DURATION of the TRANSIT is a Maximum, or a Minimum.

4107. **T**HE Calculation of the *apparent Duration* of a Transit requires no Alteration of the Place of Observation; for in one and the same Place the *Ingress* may be view'd in one Hemisphere, and the *Egress* in the other; and one shall be *accelerated* and the other *equally retarded*, by the whole Effect of Parallax. But this cannot be obtained in *one Hemisphere only*, as may easily be observed in the Diagrams.

4108. But here we cannot avail ourselves of both the *Maxima* at once, or greatest Values of the Angle $v V X = 52^{\circ} 12' 14''$, and the Side $V X = 25''$; but we can chuse a Place of Observation with either of them singly, as will appear by the following Examples.

4109. If we make this Observation with the greatest Zenith Distance of the Planet, *viz.* $Z V = 83'$ (so that $V X = 25''$) both at the *Ingress* in the *Evening*, and at the *Egress* the next Morning, 'tis plain the Meridian of the Observer must bisect the Arch of Duration; or the Complement of the Angle $Z P V$ to a Semicircle must be equal to the Arch of half Duration, which is $47^{\circ} 34' 45''$, consequently the Meridian $Z P$ of the Observer will be eastward of the Meridian of Venus $P V$ by the Angle $Z P V$

E 2

$= 132^{\circ}$

28 A CALCULATION of the DURATION

$= 132^{\circ} 25' 15''$, and therefore it will be $24^{\circ} 20' 30''$, eastward also of the Meridian of London.

4110. The Time of Half-duration $3^h 10' 19''$ (4068) taken from XII, leaves $8^h 49' 41''$ for the Time of the Evening Observation of the Ingress; and of Course, the Time of Egress in the Morning following, will be $3^h 10' 19''$.

4111. The Latitude of the Place for this Observation is thus determined in the Triangle Z P V, there are given the Side V P $= 67^{\circ} 20' 4''$, the Side X V $= 83^{\circ}$ and the Angle at P, by which we find Z P $= 22^{\circ} 10' 37''$, whose Complement is $67^{\circ} 49' 23'' =$ the Latitude sought.

4112. In the same Triangle is found the Angle Z V P $= 16^{\circ} 18' 28'' \frac{1}{2}$; to which add the Angle R V P $= 15^{\circ} 36' 26''$, the Sum is the Angle R V Z $=$ X V $\nu = 31^{\circ} 54' 54''$. This Angle being known, and the Side V X $= 25''$ will give the Side X $\nu = 13'',2$; and the Side V $\nu = 21'',2$, from which we find $\nu Y = 16''$. Then the Parallax in Longitude X $\nu = 13'',2$ gives $3' 15''$ of Time, and the Parallax in Latitude $\nu Y = 16''$ gives $3' 58''$, and the Sum of both is $7' 13''$, by which the *Ingress* is accelerated. And by the same Quantity the *Egress* will be retarded in the Morning, so that the *observed Duration* will be $14' 26''$ longer than the *central one*, and is a *Maximum*.

4113. The apparent Duration will be a *Minimum*, or the least of all, in a Situation on the South Side of Venus's Orbit, the Longitude and Latitude of which are thus determined. It is evident (Fig. 3. Plate II.) that the Ingress being retarded in the Morning, and the Egress accelerated in the Afternoon (Fig. 4.) will make it necessary that the Meridian of the Place sought should here also *bisect the Arch of Duration*, or make the Angle in the Morning Z p V $=$ Z p A ($= 47^{\circ} 34' 45''$) in the Afternoon.

4114. From whence it will follow, that the Meridian Z p of the Place will be $47^{\circ} 34' 45''$ West of the Meridian

when a Maximum and a Minimum. 29

pV passing through Venus at the Ingress, which added to $108^{\circ} 4' 45''$ (the Distance of the Meridian of *London* eastward) will give $155^{\circ} 32' 30''$, for the Situation of the Meridian of the Place, or its *Longitude, West from London*.

4115. The Latitude of the Place is thus determined for the Maximum Angle $XVv = 52^{\circ} 12' 14''$. In the Triangle ZpV there are given the Angle $ZpV = 47^{\circ} 34' 45''$, the Side $pV = 112^{\circ} 39' 56''$, and the Angle $pVZ = 36^{\circ} 35' 48''$, as we have shewn (4089.) by which we find the Side $pZ = 51^{\circ} 31' 26''$, and therefore the Latitude of the Place is $38^{\circ} 28' 33'' \frac{1}{2}$ South.

4116. Also in the same Triangle we find the *Zenith-distance* of the Planet, or Side $ZV = 75^{\circ} 41' 18''$, which will give $VX = 24'' 3$ in the Triangle XVv ; and since the Angle $XVv = 52^{\circ} 12' 14''$, we shall find the Side $Xv = 19'' 2$; and the Side $Vv = 14'' 89$, and from thence we shall have $vY = 11'' 5$.

4117. The Parallax of Longitude and Latitude together is $XY = 30'' 7$ which gives in Time of passing through it $7' 36''$, the double of which is $15' 12''$ is the Time by which the central Duration will be shortened at this Place; and is greater than that by which it will be lengthened in the North Observation (4112) by $46''$; and is therefore so much the more advantageous.

4118. As the central Ingress will in this Place be at $8h 49' 41''$ in the *Morning*, the observed Ingress will be $7' 36''$ later, or at $8h 57' 17''$, and the observed Egress will be at $3h 2' 43''$ *Afternoon*, so that the observ'd Duration will be $6h 5' 26''$, a *Minimum*.

C H A P. VIII.

Of the METHOD of finding the SUN'S PARALLAX by observing the least Distance of VENUS from the SUN. The SUPERIORITY of the METHOD of PARALLAXES; and of the PRECISION to which the SUN'S PARALLAX and DISTANCE of the EARTH and PLANETS from the SUN, may be thereby found.

4119. **A**LL that has been hitherto said relates to finding the Sun's Parallax by *parallatic Time*; but there is another Method which the Transit will furnish for that Purpose, without any Regard to Time at all; this Method consists in observing the *least Distance* of Venus from the Sun's Center, when that least Distance *S c* is a *Minimum* to an Observer on the *North Side*; and a *Maximum* when observed on the *South Side* of the Orbit of Venus.

4120. But as this is to be effected by Measurement of that Distance by Instruments properly constructed and skillfully applied; and then not by one single and direct Measure of that least Distance, but from a Result of many Measures taken from various Distances of the Centers of the Sun and Venus whilst she is traversing his Disk, or from many Observations made on the Differences of the *right Ascension* of those Centers during the Passage over the Disk; I say, upon these Accounts, it will prove too critical and difficult an Affair for every common Observer to undertake.

4121. In a former Part of these Institutes (3540, &c.) I have given the Construction of an *Helioſtata* or *Celestial Clock*, by which the Image of the Sun is rendered *quiescent*, or without Motion. By Means of a common reflecting Telescope this Image may be made 12, 15, or 18 Inches Diameter; so that every single Second of a Degree becomes distinctly visible to the Eye, and the Whole performed
without

the SUN'S PARALLAX may be found. 31

without any Difficulty or Trouble. Those who are judicious in these Matters need not to be informed with how great Facility any Measures of the Distances of Venus from the Sun's Center may be taken during the Time of the Transit: This every one has acknowledged, to whom I have shewn the Effects of such o'Clock.

4122. Indeed it would be more worth while to insist upon this Method of measuring the least Distance of the Centers, were it not in itself so deficient from answering the Purpose intended by it, *viz.* for estimating the Parallax of the Sun; for in the Instances we have given the Differences of the *central and observed least Distances*, when reckoned in Defect and Excess, do in the Whole, make no more than 36"; and suppose they amounted to 40", you could only find the Parallax within a 40th Part of the Whole; which would give but little Satisfaction as it would leave so great Uncertainty in the Distance of the Sun. For Example, suppose the Sun's Parallax = 10", and the Semidiameter of the Earth = 4000 Miles, then the Distance of the Sun would be 82500000 Miles; a 40th Part of this Number is 2062500; so that more than *two Million of Miles* would depend upon taking that least Distance true to one Second of a Degree.

4123. The Method therefore, which principally, if not solely, deserves Attention, is that by *parallatic Time*, both on Account of its Facility and the greatest Precision we can ever hope to arrive at. As to the former, (to use Dr. HALLEY's Words) this great Problem is solved without much Labour, — and without any other Instruments than Telescopes, and good common Clocks; and without any other Qualifications in the Observer than Fidelity and Diligence, with a moderate Skill in Astronomy.

4124. Then as to the Precision, or Degree of Truth in the Determination, we have shewn there is 15' 12" in Excess (4117), and 14' 26" in Defect (4112), with regard to the Duration of the Transit from that observed at the Earth's

Earth's Center; so that by this Species of Observation, we have a Scope or Extent of $29' 38''$, or of $1778''$ to estimate the Sun's Parallax by.

4125. But if we take the Differences of parallatic Time at the *Ingresses*, and *Egresses*, we shall have still a larger Number of Seconds, *viz.* $31' 6'' = 1866''$. And the aforesaid Dr. HALLEY has assured us, that he could very easily observe the internal Contacts of the Planet Mercury and the Sun, *without an Error of a single Second of Time*; and from thence he concludes, that, in a Transit of Venus, these Contacts may be observed so accurately, that the *Sun's Parallax may thereby be determined to a small Part of a Second*; and in another Place he affirms, the *said Parallax may be known within a 500th Part of the Whole*. And he was too great a Judge of such Subjects to leave the least Doubt of the Truth of what he has asserted.

4126. But after all, these Sums of Differences found at two different Places, depend upon precarious Principles, *viz.* the ascertaining the *true Longitude of those Places*; because *every Minute's Difference in Longitude will make a Difference of four Seconds of Time*. Therefore I think the surest Way will be to depend upon those Observations of the *Duration* that are made in particular Places on the North, and on the South Side of the Orbit; where the Difference from the Central Duration will be found sufficiently large, and very safe from Error; for $15' 12''$ give $912''$, by which if you divide 82500000 , the Quotient 90460 will be the *Number of Miles corresponding to each Second of Time*.

4127. Whence it appears, that if the Parallax of the Sun could be really determined to the *900th Part of the Whole*, we should still be at the *Uncertainty of 90000 Miles in the Distance of the Sun from the Earth*. And yet this is perhaps a nearer Approach to it than we may ever be able to make; since in order to this, the Observation in all its Circumstances, must be *just to a single Second of Time*.

the SUN'S PARALLAX may be found. 33

4128. Since all Effects are proportional to their Causes, it will follow, that the calculated parallatic Difference of $15' 12''$ is to the assumed solar Parallax of $10''$ as the observed parallatic Difference of Duration at the Time of the Transit is to the Parallax of the Sun, as near the Truth as such an Observation can give it. Thus for Example, suppose at the Place mentioned (4117) the Time of the Duration of the Transit to be found to be $6^h 34' 19''$; this is greater than the central Duration $6^h 20' 38''$, by $13' 40'' = 821''$; therefore as $15' 12'' : 13' 40'' :: 10'' : 9'' = \text{Sun's Parallax}$, as truly as it can be obtained by the very best Observation.

4129. Again, suppose the Duration of the Passage be observed at the same Place to be $6^h 36' 36'' \frac{1}{2}$, this exceeds the central Duration by $15' 57'' \frac{1}{2} = 957 \frac{1}{2}$, and therefore you say, as $912'' : 957 \frac{1}{2} :: 10'' : 10'' \frac{1}{2}$; so the Sun's Parallax will in such a Case be 10 Seconds and a Half. But if the observed Duration there be just $6^h 35' 50''$, which is $15' 12''$ longer than the central Duration, then the Parallax of the Sun will be known to be 10 Seconds.

4130. In the Triangle A B E (Fig. 4. Plate I.) since A E continues the same, it is plain the Side A B will be greater or lesser in the *inverse Ratio* of the Angle A B E, that is, the Distance of the Sun A B from the Earth E, will be *greater or less* than 82500000 Miles, in proportion as the solar Parallax A B E is found by Observation to be *less or greater than 10 Seconds*.

4131. Now the proportional Distances of the Planets from the Sun are known by the *periodical* Times resulting from the Law of Gravity; and are thus stated, viz.

For ♀	♀	☉	♂	♃	♄
38718	72333	100000	152369	520096	954006.

4132. Therefore, if the Distance of one could be once known, those of the other Planets will be known by the Rule of Proportion. Suppose the solar Parallax be found $= 10''$, the Distance of the Earth will 82500000 Miles; then say,

34 *The GEOGRAPHY of a TRANSIT*

as 100000 : 825000000 :: 38710 : 32000000 = Distance of Mercury from the Sun in Miles; and so for the rest.

4133. Not only the *Distances* of the Planets, but also their *Magnitudes* are known by the Parallax of the Sun. For since the *real Diameters* of Bodies are in the same Ratio of their *apparent Diameters*, or Semidiameters, and that of the Sun (at a Mean) being $15' 30'' = 930''$, and that of the Earth = (the *Sun's Parallax*) $10''$; therefore the Sun's Diameter is to the Earth's Diameter as 930 to 10, or as 93 to 1; and so the Sun's Magnitude is to that of the Earth, as 804357 to 1.

4134. By the last Transit 1761, the apparent Diameter of Venus was found to be $58''$; therefore, if Venus were viewed at the Distance of the Sun, her apparent Diameter would be less in the Proportion of 100000 to 27667, that is, it would there be only $16''$, but the Earth's Diameter at that Distance is $20''$ (*viz.* double the Sun's Parallax) therefore, the true Diameter of the Earth is to that of Venus as 20 to 16, or as 5 to 4; and, of course, the Magnitude of the Earth is to that Venus, as 125 to 64, almost twice as large.

4135. Hence it follows, that the *Magnitudes* of the Sun and Planets will be greater or less in Comparison of the Earth, as the Sun's Parallax at the next Transit shall be found less or greater than 10 Seconds. Therefore the final Determination of the *Dimensions* of the PLANETARY SYSTEM depends on that Discovery.

C H A P. IX.

The GEOGRAPHY of a TRANSIT illustrated by a PRAXIS on the Terrestrial GLOBE, exhibiting all the Phænomena of the ensuing TRANSIT of VENUS to the various INHABITANTS of the EARTH.

4136. **T**HE GEOGRAPHY of a TRANSIT consists in a general Representation of all its Phases, in respect of the different Situation of the Inhabitants of the Earth, by the *terrestrial Globe*; and to point out and distinguish all those particular Places upon the Surface of the Earth, or Ocean, where the *Beginning, End, and Duration of the Transit*, may be observed to the greatest Advantage possible, for the Discovery of the *solar Parallax*, and, in consequence thereof, the *Dimensions of the SOLAR SYSTEM*.

4137. In order to this curious and delightful Praxis, every Thing previously necessary has been already calculated; and to see the Reason thereof, one Thing only is to be considered, *viz.* that as the Sun enlightens one half of the Globe constantly, it is therefore always in the *Vertex or Zenith of that enlightened Hemisphere*; and that the *Circle of Illumination* which terminates the light and dark Hemispheres, may be considered as a *solar Horizon*; and lastly, that the Meridian in which the Sun is, is the *Eastern Horizon* in which he appears to be *rising* to all the Inhabitants of the Earth under the *Western Semicircle of Illumination*; and the same Meridian is the *Western Horizon*, in which he appears to be *setting* to all the Parts of the Earth under the *Eastern Semicircle of Illumination*.

4138. Therefore, to rectify the Globe for the ensuing Transit of 1769, *June 3d*, the North Pole of the Globe must be elevated to the Sun's Declination, which is $22^{\circ} 27' 15''$; (4062) then will this Point of the Sun's Declination on the South Part of the Brass Meridian truly represent

36 *The* GEOGRAPHY of a TRANSIT

the *Sun's Place* during the Time of the Transit, as being the *Zenith* of the illumined Hemisphere.

4139. Then, since the Time of the *beginning* of the Transit is computed for LONDON, and is found to be at 7^h 12' 19" in the Evening; (4069) therefore bring the Meridian of *London* to coincide with the general (or Brass) Meridian, and holding the Globe fast in that Position, set the *Hour-Index* to vii^h 12' at Night; after this, turn the Globe Eastward till the said Index points to the upper XII, and there let it rest.

4140. Then it will be *Noon* to all those Places which lie under the upper Semicircle of the general Meridian; and of Course they will see the beginning of the Transit, or first Contact of Venus, with the Limb of the Sun upon their Meridian, or at 12 at Noon. These Countries you'll observe are only the *Western Parts of North America*, the western Parts of *Mexico*, by the Cape of *California*, and then over the vast *Western Ocean* to 67° 33' of South Latitude. The Sun will be vertical at that Time to a Place near Cape *St. Lucas*, the South Point of *California*.

4141. To all those Places which lie under the Western Semicircle of the Horizon of the Globe, the Sun will rise with Venus in Contact on his Eastern Limb. These are the *White Sea*, the most Northern Parts of *Siberia*, almost through the Middle of *Great Tartary*, the Northern Parts of *Japan*, and many of the Islands in the *Great South Sea*.

4142. Lastly, all those Places under the Eastern Edge of the Horizon will see the beginning of the Transit at *Sunset*; viz. from the *White Sea*, the Northern Parts of *Russia*, South East Parts of *Denmark*, the Southern Parts of the *Baltic Sea*, *Poland*, North West Parts of *Germany*, South East Parts of *France*, the Eastern Coast of *Spain*, the middle Parts of *Morocco*, and all the Kingdoms upon the Western Coasts of *AFRICA*, and along the *Great Ocean* towards *South America*.

4143. From what has been said, 'tis evident, that all Parts of the Earth *between the Western Horizon and the Meridian*, see the beginning of the Transit in the *Morning or Forenoon*; and those *between the Eastern Horizon and Meridian*, see it in the *Afternoon*; at different Hours according to their different Longitude or Distance from the Meridian.

4144. After the same Manner you proceed for the *middle of the Transit*, which will be at $10^h 22' \frac{1}{2}$ at Night; therefore bring LONDON to the Meridian, where hold the Globe fast, and set the Hour Hand to $10^h 22' \frac{1}{2}$, and then turn it till the Index comes to the upper XII. Then, in the Western Horizon, you see all the Countries who observe the Planet in the middle of her Path upon the Sun's Disk at *his rising*; under the Meridian are all those Places who have that Phase at Noon Day; and in the *Eastern Horizon* are all those which see the same at *Sun-set*. Also in this Position of the Globe you have in one View all the Places who view the middle of the Transit in the *Forenoon*, and in the *Afternoon*, at every Hour.

4145. The *End* of the TRANSIT will be at $1^h 33'$ in the Morning of *June 4th*; therefore placing LONDON under the Meridian, and the Hour Hand at the same Time to $1^h 33'$, turn the Globe eastwards till the Index comes to 12 at Noon, and there let it rest. Then in the Western Horizon are all the Places who see the *Sun-rising* at the End of the Transit; those under the Meridian see the last Contact at *Noon*; and the same Phase appears at *Sun-set* to all who are in the Eastern Semicircle of the Horizon.

4146. From what has been said it will follow, that if when the Globe is rectified for the beginning of the Transit as directed (4139) you draw a Circle quite round the Globe by the interior Edge of the Horizon with a piece of Chalk or Crayon, and afterwards such another Circle, when the Globe is rectified for the End of the Transit (as in the last) then will those two great Circles intersect each

38 *The GEOGRAPHY of a TRANSIT*

other in Points $31^{\circ} 29'$ distant from the Poles of the World, and divide the Surface of the Globe into four unequal Parts, or Gores.

4147. Then in that Gore, which is contained between the two eastern Semicircles, which shew the Sun *setting* at the Beginning and End of the Transit, you will see those Places *where the beginning of the Transit will be seen, but not the End*; viz. the Eastern Parts of *North America*, all the *West Indies*, and *South America*; the Western Parts of *Europe* to the Latitude of 65 Degrees; the Western Parts of *Africa*, and all the *Atlantic Ocean*.

4148. In the Quarter or Gore opposite to this, you observe those Places only *who see the End of the Transit but not the Beginning of it*; because at the Beginning, this Gore was wholly in the nether or dark Hemisphere. The Countries herein comprehended are *Russia*, and the greatest Part of the Continent and Islands of *Asia*, *New Holland*, *Zealand*, &c.

4149. In that Quarter bounded by the two Semicircles, of which one shews the Sun *rising* at the beginning of the Transit, and the other shews him *setting* at the End of it, are contained all those Parts of the World who see the *Whole of the Transit*; because they are in the illumined Hemisphere, or above the Horizon, the whole Duration of the Transit. This Quarter of the Globe is almost wholly possessed by the *Pacific Ocean*, or *Great South Sea*, and its numerous *Islands*. In the Northern Parts of it are contained the most Part of *Great Tartary*, *Nova Zembla*, all *Lapland*, *Baffin's Bay*, *Hudson's Bay*, and all the unknown Part of *North-West America*.

4150. In the Quarter opposite to this, you find the Situation of all those Countries whose Inhabitants see *nothing at all of the Transit*. Such are the *midland Parts of Europe*, all the *Mediterranean*, all the *Arabia's*, and part of *Persia*, and nearly the whole *Continent of Africa*, with the *vast Ocean* to the South thereof; for these Parts will be under the
Horizon

Horizon during the whole Passage of Venus over the Sun's Disk.

4151. Since the Declination of the Sun $22^{\circ} 27'$ is so near the greatest Declination $23^{\circ} 30'$, it will follow, that a Tract about the North Pole, nearly equal in Extent to the whole *Frigid Zone*, will at that Time of the Year have the Sun constantly above the Horizon; the Inhabitants of one Part of this Zone will have a Motion in the same Direction of the Planet Venus, *viz.* from East to West; and this will cause the Duration of the Transit to be longer to them than the central one; and the Inhabitants of the opposite Parts of the same Parallel, as they move in a contrary Direction, will observe the Duration to be shorter than that seen from the Earth's Center.

4152. But as we have already determined *six different Places* where the Differences by parallatic Time are the *greatest and least possible*, with regard to the *Ingress, Egress, or whole Duration*, we shall next point out the Situation of those Places upon the Surface of the Globe by means of their given Longitudes and Latitudes.

4153. We have shewn that the Place where the Ingress is most of all accelerated, is in North Latitude $51^{\circ} 28'$, and Longitude $6'$ East from *London*; and that points out the City of LONDON itself for the best Situation to observe that Phase (4090).

4154. The retarded Ingress is a *Maximum* in South Latitude $43^{\circ} 30'$, and Longitude West from *London* $162^{\circ} 49'$; this Longitude brought to the Brass Meridian will give the Point under the $43^{\circ} 30'$ for the Scite of the Place required, which is in the southern Part of the *Pacific Ocean*. (4091).

4155. The Place where the retarded Egress is seen to the best Advantage has Latitude 23° North, and Longitude East of *London* $68^{\circ} 34'$ (4097), therefore bring this Longitude to the general Meridian, and under the 23° of Latitude you will find the Place to be a little above *Surat* on the Gulph of *Cambay*, near the *Tropic of Cancer*.

40 *The GEOGRAPHY of a TRANSIT*

4156. And the Place where this Phase is most of all accelerated is in *South Latitude* $17^{\circ} 16'$, and $128^{\circ} 44'$ West of *London*. (4103). This will give the Situation of the Observer in the *Pacific Ocean*.

4157. Then again, we have shewn (4112) that the Place where the Duration will be observed a Maximum is in *Latitude* $67^{\circ} 49'$ North, and *Longitude* $24^{\circ} 20'$ East of *London*; this therefore will be found a little above *Tornea*, in *Swedish Lapland*.

4158. And lastly, the Place where the Time of the Transit will be observed a Minimum, is in *South Latitude* $38^{\circ} 29'$, and *West Longitude* $155^{\circ} 32'$; which determines the Place in the *Pacific Ocean*, or *Great South Sea*. (4118).

4159. It is further to be remarked, that tho' the six Places above determined are indeed the best that can be, yet a Distance of many Leagues round either, will make but a trifling Difference in parallatic Time; and therefore the Observer may look upon these Places only as *Indices* of what Parts of the World will be most proper for his Station and Observations. Thus 50, 80, or 100 Miles round *LONDON*, will be nearly of the same Advantage as the City itself. All the *Mogul's* Empire about *Surat* and *Camboy* will do for Observations there; and over all *Norway* and *Lapland*, the *Northern Observations* will be equally good.

4160. On the other Side the Line we have the three Stations near the same Part of the *Great Ocean*, and all beyond the Islands so numerous in lesser Latitudes, and other Parts of the same Ocean, which nevertheless will be found good Situations for those *Southern Observations*.

4161. As we have been so large and particular in the *geographical Praxis* by the Globe, it will be quite unnecessary to give a *geographical Map of the Transit*, as it is much more difficult and less perspicuous and pleasant to study the Geography of the Transit that Way, than by a *Transit Globe*, with proper Circles drawn upon it, and all the

the Divisions or Gores properly distinguished by different Colours where the various and different Phases and Quantities of the Transit will be observed over the Face of the whole Earth.

4162. I cannot leave this Subject without remarking an Error in all the Maps of the Transit I have seen, viz. that it is asserted by their Authors, *that to all the Space contained between the two Semicircles mentioned in (4149) the Whole of the Transit will be visible; and to all the Space of the opposite Gore (in 4150), the Whole of the Transit will be invisible.* Now, that this is not true will appear by rectifying the Globe for the *middle of the Transit* as directed in (4144), and drawing a Circle upon the Globe quite round the Horizon, for then it will be seen that the Gore of perpetual Apparition (4149), is not wholly in the upper and illuminated Hemisphere, but a Part at the *North End* will be cut off by this Circle, and consequently must be below the Horizon at the middle of the Transit, and to them therefore that Phase will be invisible.

4163. On the other Hand, a Part of the Gore mentioned in (4150) will be in the enlightened Hemisphere at the middle of the Transit, and therefore to some Part of this Space, some Part of the Transit will be visible. This re-scinded Part of either Gore is in Form of an *Isosceles Triangle* whose Altitude is 9 Degrees, and its Base about 19° ; and of course not so very inconsiderable as to be wholly neglected in an accurate Explication of the GEOGRAPHY of a TRANSIT. But we shall further consider this Subject in a Chapter on the *Transit-Globe*.

C H A P. X.

The NATURE of a TRANSIT and its various Phenomena explained, by an Orthographical PROJECTION of the EARTH, SUN, and Planet, upon a Perspective PLANE, and a general COMPUTATION of parallatic TIME from thence.

4164. **T**HERE is another Method of treating a Transit, which can by no Means be omitted in a System of Institutions of this Kind; the Knowledge of which no ingenious Student in Astronomy can be supposed to dispense with; which is in itself most elegant and geometrical; and lastly, a Method which gives the general Rationale of all others; it consists in an *Orthographical Projection* of the Earth, the Sun, and Planet, upon a perspective Plane passing through the Center of the Planet, and perpendicular to the Axis of a Cone of Rays, from a Point of Sight in the Center of the Sun.

4165. The Principles of this Projection, applied to *Solar and Lunar Eclipses*, have been already largely explained in these Institutes (1721, &c.); they are equally necessary for a true Representation of a Transit *in Plano*, as will appear from a due Consideration of Fig. 4. Plate I. wherein the Diameters, or rather Semidiameters, of the Sun, the Earth, and the Planet, are evidently as the Angles under which they appear as viewed from the Earth, Sun, or Planet. It has also been shewn, that the Parallax BC of the Planet V by two Observers, one at the Earth's Center E , and the other upon its Surface at A , will be the Angle BVC , or AEC ; but the same Parallax BC will be seen by the Observer at A only, under the Angle BAC .

4166. But this Angle BAC depends upon the Planet's Distance from the Sun, as it increaseth and decreaseth therewith; also it is well known, that the Perspective of any Object

Object upon a Plane, will increase and decrease with the Distance of that Plane from the Eye. From all which it follows, that if the Eye be supposed to be placed in the Center of the Sun at S, and a perspective Plane at the Orbit of the Planet P V, perpendicular to the visual Line S P E, then a Circle described upon the Center P of such a Plane, with the Radius of 945 equal Parts or Seconds, will represent the Disk of the Sun, for so many are contained in his Semidiameter S D = $15' 45''$, as viewed from the Earth at E. (Fig. 4. Plate I.)

4167. Again, to an Eye at the Sun in B, the perspective of the Earth's Semidiameter A E will be equal to Q V upon the Plane. But A E is to Q V, as the Parallax of the Planet A V E to the Parallax of the Planet from the Sun Q A V. That is, in our present Case of the Planet *Venus*, we have A E : Q V :: $35'' : 25''$ (4078), consequently the Semidiameter of the Sun and Earth upon that perspective Plane, will be as 945 to 25.

4168. Therefore, with a Radius S E = $945''$ (taken from a Scale of equal Parts) describe on the Center S the Arch E F, to represent a Part of the Limb of the solar Disk; and upon the same Center S with a Radius S a = $25''$ (from the same Scale) describe the small Circle a f b g, to represent the Perspective of the Earth's illumined Hemisphere; then will a b represent that Meridian in which the Sun is at S, a g b is the eastern Semicircle of Illumination, and a f b is the western Half. (See Fig. 1. Plate III.)

4169. Let the Line C V be half the Transit Line A V in the Diagrams of Plate II; and because the Conjunction at D happens at 10 o'Clock (4057), therefore if you take $242''$ from the Scale, and placing one Foot of the Compasses in D, cross the said Line with the other Point towards the right Hand, it will give the Hour of IX; from thence you set the same Distance to VIII. and then to VII. and thus

the Path of Venus is divided into Hours, each of which is subdivided into Minutes sufficiently distinct to the Eye; so that the Place of Venus is seen for *every Minute of her Passage*.

4170. Let SC be perpendicular to the Path, as in the other Figures, and let V, E, G , be the Points in which the Planet is at the external Contact, central Ingress, and interior Contact; the Time of the former is $7^h 12' 19''$ (4069), and of the latter, $7^h 30' 55''$ (4070), as seen from the Earth's Center S . And taking the Extent from S to V in the Compasses, place one Point in V , and with the other describe a small Arch ih ; then will all the Inhabitants of the illumined Disk over which the Arch ih passes, see the beginning of the Ingress at the same time it would be viewed from the Earth's Center. In the same Manner you proceed for the central Ingress, and internal Contact at E and G .

4171. The same Extent or Line SV being taken in the Compasses, one Foot of which being placed on any particular Point of the Earth's enlightened Surface, the other will fall upon the Minute of Time in the Path V , at which the external Contact will happen to the Inhabitants of that Place.

4172. It is evident from this Operation, that to all Parts of the Earth's Disk which lie beyond the Arch ch , the Time of the beginning of the Transit will be *later* than the central Time; and latest of all at the remotest Point r of that farther Hemisphere.

4173. On the other Hand, one Foot of the Compasses being placed upon any Part of the near Hemisphere, the other will fall upon that Minute in the Transit Line at which the beginning of the Transit happens in that Place, which of course will be *before the Time at the Center*, or in the Arch ih ; and the most early beginning will be to those who live at o , in the Part nearly opposite to r .

4174. If the same Construction be made for the Planet's Egresa from the solar Disk, then the Moments of the Phases of the beginning, central, and total Egresa, will, in like Manner, be seen, or found for all Places upon the Earth's Disk (*a f b g*).

4175. In all that has been hitherto said, the Reader will observe, that the Triangles *SCV*, &c. and all the Sides and Angles thereof, are of the same Quantity as those lesser ones mentioned (4065), but here the Projection is large enough to give a distinct Idea of the Manner in which the Moments of Time for the various Phases of the Transit are to be ascertained for the several Parts of the Earth's Surface, where it can in any Part be seen.

4176. If the projected Disk of the Earth (*a f b g*) be three Inches in Diameter, (as in Fig. 4.) the Line *SV* will be 4 Feet 10 Inches; then that Distance taken in the *Beam Compass*, and the Point placed in *s* and *r*, the Pencil will mark the Minute on each Side the Point *V* in the Path, which denote the earliest and latest Moments of the beginning of this Sort of Eclipse; then afterwards place the Point of the Compass upon each Minute between these two successively, and upon them, as Centers, strike circular Arches upon the Earth's Surface with the Pencil; these Arches will divide the then enlightened Surface in such Manner as to shew all the Places where the beginning of the Transit differs by one Minute of Time. And thus you divide the Earth's Surface for the Minutes of the Difference of Time at the end of the Transit.

4017. From what has been said, it will be easy for the young Student in Astronomy to understand how the same Minute Arches may be drawn upon the Surface of an artificial Globe itself, of 3, 6, or 9 Inches Diameter, by placing it in a circular Hole of a board, or Plane, with half a Meridian to move upon the end of the describent Line, and to be set at right Angles to it.

4178. It now remains to assign the Seat or Perspective of any proposed Place upon the Earth's projected Disk; for this Purpose the Latitude and Longitude from the Meridian *ab* must be known; as also the Position of the Plane of the Meridian passing through the Place, and the Elevation thereof above the primitive Circle, or Plane of Projection. To find the two last Requisites proceed thus; let *AB* be the Meridian of the enlightened Hemisphere passing through the Sun *S* in the Center; *BP* the Elevation of the North Pole *P*, stereographically projected; and suppose *CPE D* the Meridian of the Place making the Angle *EPS* or *CPB* = 35° , or given Longitude, from the first Meridian *AB*. Then in the right-angled spherical Triangle *BPC*, there are known the Side *BP* = $22^\circ 27'$ the Sun's Declination, and the Angle *CPB* = 35° ; by which you find the Side *BC* = *AD* = 15° , and the Angle *BCP* = 58° . (See Fig. 2. Plate III.)

4179. Then take *SB* in the Compasses and make it a Radius on the Line of Sines upon the Sector; and then take the parallel Distance from 32 to 32, (the Complement of 58°) and set it from *S* to *L*; so will *SL* be the semi-conjugate Axis, and *SC* the semi-transverse of the elliptic Meridian passing thro' all Places of 35° Longitude from the Meridian *AB*; and may be drawn as directed (785.)

4180. Let the Latitude of the Place be 40° ; the elliptic Parallel of that Latitude is therefore next to be drawn. In order to do that, you proceed thus, by the orthographical Projection; let *EPQp* be the solar Meridian of the illumined Hemisphere placed directly before the Eye (Fig. 3. Plate III.) its Diameter *AB*; Center *S*; *EQ* the Equator; *Pp* the Axis; *CD* the Diameter of the Parallel of 40° = *EC*, and its Center, *N*; also *cd* is the same Parallel of South Latitude, and its Center *n*; and ☉ the Sun in the Zenith.

4181.

4181. Then right Lines perpendicular to $A B$, passing thro' the Points $P, N, C, *, n c$ will project those Points into the Points O, R, T, S, r, t , in the Axis $A B$; and the Radius $S B$ set from 90 to 90 in the Sines upon the Sector, you will have the parallel Distance from $67^{\circ} 33'$ upon the same Sines equal to $S O$, for the projected Pole O .

4182. The projected Center of the parallel is R , which is therefore the Center of the Ellipsis to be drawn; the longest Diameter of the Ellipsis will be equal to the Diameter $C D$ of the Parallel, or the Semi-transverse is equal to $C N$.

4183. The Projection of the Point C of the Parallel in the Meridian is T , and its Distance from the Center S is $S T$, equal to the parallel Distance of the Sines $17^{\circ} 33'$ on the sectoral Lines; that is, $S T$ is the Sine of the Arch $* C = E C - E *$, or Difference between the Latitude of the Parallel and the Sun's Declination. But $S t$ is the Sine of their Sum, viz. $c E + E *$, for the same Parallel in South Latitude. From hence it is evident, that $T R (= t r)$ is the semi-conjugate Axis of the required Ellipsis, and therefore having both the Axes, the Ellipsis is drawn as directed in (785)

4184. Therefore (in Fig. 4.) let $A B$ be the Meridian of XII. on the illumined Disk of the Earth, and let $K M L$ be the Meridian of the given Place in 35° West Longitude of $A B$; (4179.) also let $C T D$ be the Parallel of 40° the Latitude of the Place proposed, which is now determined upon the Disk in the Point of Intersection M of the said Meridian and Parallel.

4185. Then taking $S V$ in the Compasses (Fig. 1.) and placing one Foot in the Point m upon the Earth's Disk, extend the other to the beginning of the Path towards V , and it will fall upon the Minute of Time at which the Contact at V will happen to a Spectator on the Earth in Latitude 40° and West Longitude 35° . If you take the
Line

Line S G in the Compasses and apply it as before, it will shew the Minute when the internal Contact will happen at the same Place.

4186. If a Calculation to determine the Point M be more eligible, you proceed thus: As Radius is to the Sine of the Co-Latitude C O, so is F S to C R. Also C R is to M V as Radius to the Sine of the Longitude $40^\circ = \text{Arch M T}$; therefore by Composition of Ratios, we

$$CO \times MT \times FS$$

have $\frac{R^2}{R^2} = M V$.

4187. Again, Radius is to the Sine of C M $= 55^\circ$, as R T is to W M $= R V$; and S R $- R V = S V$; then having the Length of S V and V M, the Point M is thereby ascertained upon the Disk.

For Example, the Sine of C O $= 50^\circ = 9.884254$

Sine of M T $= 35^\circ = 9.758591$

F S $= 25 = 1.397940$

From the Product 21.040785

Subtract $R^2 = 20$.

There remains the Value of M V $= 10'',98 = 1.040785$

4188. Then by the Analogies above (wrought upon the Sliding Rule) you will find S R $= 16''$; and Radius is to the Sine of the Arch O C (Fig. 3.) as F S is to S T $= 7'',6$. Then S R $- S T = 8'',4 = T R$; and lastly $R : CM = 55^\circ : : T R : M W = R V = 6'',88$. Then S R $- R V = S V = 9'',12$. Having thus found V M and V S in Numbers, the Point M is given upon the Disk by the plain Scale and Compasses.

CHAP.

C H A P. XI.

Of the ANALOGY and CORRESPONDENCE between the Orthographical PROJECTION, the METHOD by DIAGRAMS, and the PRAXIS by the GLOBE, in regard to the various PHASES of a TRANSIT.

4189. **I** Shall now point out the Analogy and Correspondence there is between this orthographical PROJECTION and the Method by DIAGRAMS in Plate II. and the Praxis by the *Terrestrial GLOBE* in Chap. IX. In order that this may be more evidently made out, it must be first observed, that in such a Projection as the present one, the Vertex or Zenith of the Place under the Sun is the Center thereof, or the Point S, and that all the vertical Circles S Z being at Right Angles to the Plane of the Primitive, are projected into right Lines S Z, and that the Zenith Distance of any Place M is S M; and whose Measure is on the Line of Sines, S Z being Radius.

4190. Therefore since Radius is to the horizontal Parallax S Z, as the Sine of the Zenith Distance is to the Parallax S M pertaining to the Place M, (4021). It is plain that the Line S M in this Projection corresponds to the Line V X in the Diagrams; and again, let the Diameter P Q be the Axis of Venus's Orbit upon the Disk, and from the Point M draw M v perpendicular to it, then will the Parallax of Altitude M S be thereby resolved into two others, viz. S v, which is the Parallax of Latitude from the Orbit, and the other is M v parallel to the Orbit, and therefore is the Parallax of Longitude in the apparent Path. Consequently S v and M v do here answer to V v and X v in those Diagrams, therefore the Triangle S v M and V v X are analogous, or the same Thing in different Views.

4191. If N represent a Place in the same Parallel of Latitude, and in the same Longitude, but on the other Side or Eastward of the Meridian AB , then drawing the Perpendicular Ns , the Line Ns will be the Parallax in the Orbit; and the Line Ss is the Parallax of Latitude; if Ns be continued it will cut Sy in w ; and the Triangle SwN will answer to the parallatic Triangle VYX in the Diagrams, and sw is the efficacious Part of the Parallax of the Latitude Ss , because the Angle $NSs = XVv$; and $Ss w$, or $PSy = VvY$.

4192. If M be the Site of a Place in the Western Hemisphere where the Transit begins, and N that in the Eastern Hemisphere where it ends, and $TM = TN$; then Sv will be the Parallax of Latitude at the beginning, and Ss that at the end; and so the apparent Path mn of Venus to that Place will be inclined to the true or central Path MN in the Diagrams, and not parallel, as there represented.

4193. Draw ase at right Angles to SP , and let E be a Place near the Western Horizon in the Parallel of the Sun's Declination where the Transit appears to begin, and S its Place when it ends; then draw Eb perpendicular to Se , and it will be the Latitude or Distance of the apparent Path, and the true one at the beginning of the Transit; but at the End, they both coincide. And what is more, all Places (circumstanced as above) between North Latitude E and South Latitude a , will be on the North Side of the Orbit at the beginning, and on the South Side at the end of the Transit; and so the apparent Path mn , (in the Diagrams) will intersect the central one MN , or lie across it.

4194. It is further evident, that to every Place N in the eastern Hemisphere PQ , the Parallax of Longitude Ns conspiring with the Motion of Venus, will accelerate every Phase of the Transit; and if the Place be on the North Side or Hemisphere aP , then also the Parallel of Latitude

sw will be directed the same Way, so the Phase will be accelerated by the Sum Nw of both Parallaxes, consentaneous to (4051.)

4195. But if the Place be on the southern Side of the Orbit, or Hemisphere, aQe , then the Parallax of Latitude has a contrary Tendency to that of Longitude, and the Phase of the Transit is accelerated by their Difference so long as that is positive, that is, so long as the Parallax of Longitude is greater than that of Latitude.

4126. For since after you pass the Orbit of Venus aSe , the Parallax of Latitude becomes negative and increasing, while that of Longitude constantly decreases, there must be some Point N in every Secondary PNQ , where the negative Parallax of Latitude will be equal to the Parallel of Longitude Ng . And any Place to the Southward of that, or towards Q , will have the Phase retarded by the said Differences of Parallaxes.

4197. In the Points P and Q , where the Longitude vanishes, and where the Effect of the Parallax of Latitude is greatest of all, the Value of this Parallax will be expressed by the Line QY , or Tangent of the Angle QSY compared with Radius QS . But the greatest Effect of the Parallax of Longitude is at a , or e , and is the horizontal Parallax itself, viz. Se or $Se = 25''.14$, (4077) therefore the greatest Effect of the Parallax of Longitude is to that of Latitude, as Se or QS to QY , or as Radius to the Tangent of the Angle $QSY = 37^\circ 48'$, in the present Case.

4198. An Observer at any Place M in the Western Hemisphere PaQ , and on the North Side of the Orbit ase , will observe the Phases of the Transit accelerated, but it will be only by the Difference of Parallaxes; for the Parallax of Longitude Mv retards, and that of Latitude vu accelerates them, and the Difference Mu is positive; but that vanishes when the Place M is in

the Line SY ; [and afterwards, in all the Hemisphere hri , both the Parallaxes conspire to retard the Phenomena of the Transit. But in the Parts of the Disk between yS and aS , the Retardation is effected by the Differences of the two Parallaxes, that of Longitude being greatest and negative. In all Parts between aS and SQ the Sum of the Parallaxes retard the Times. Lastly, from SQ to SY , they are again retarded by the Difference; since there the Parallax of Latitude prevails over that of Longitude, and is contrary to the Motion of Venus.

4199. The Line ib (in Fig. 1.) which divides the Surface of the Earth's Disk into the two Parts iob and irb , in the present Construction of Fig. 4. coincides with the Line yY ; and the Angle oSP is the Maximum Angle, and equal to the Angle ZVR , or XVv , in the Diagrams of Plate II. Also the Angle iSP is equal to the Angle vVY there. If therefore the Line ok be drawn parallel to ae , it will cut PS in l , and the Part ol will be the horizontal Parallax of Longitude, and lk that resulting from the Parallax of Latitude Sl ; both accelerating the beginning of the Transit in the greatest Degree possible.

4200. In order to find the Quantity of this greatest Acceleration (and that of any other) the two Scales AB and CD may be made, as in Fig. 5. The first AB is equal to the Radius of the Disk, and is therefore divided into $25''$, 14 being equal to the horizontal Parallax of Venus from the Sun (4077); upon this the *Number of Seconds of a Degree* in any Parallax is measured. And the Scale CD is divided into *six Minutes of Time*, and is equal to $24,2$ Seconds of Motion in the Scale AB , as is evident from (4067).

4201. Thus for Example, take ol in the Compasses, and measure it upon the Scale AB , and it will be equal to $20''$ of Motion; upon the Scale CD it will be equal to $5'$ of Time in which that Motion is performed. Then lk
(the

(the Effect of Latitude) measured upon A B is near $12''$ of Motion; upon C D it is near $3'$ of Time; so that both these *Maxima* together make almost $32''$ of Motion, or nearly $8'$ of Time; which is the greatest Quantity by which the beginning of the Transit can be accelerated. By considering the Position of the Meridian O o, and that of the Point o, they will be found near to those of *London*, as will appear by rectifying the Globe as taught in (1453).

4202. Hence also it appears that the greatest Effect of Latitude by which the Phase can be accelerated is P y, and equal to Q Y by which it will be retarded; and if measured upon A B will be found $19'' \frac{1}{2}$; and upon C D it will give about $4' 50''$ of Time.

4203. I shall only remark further, that in the Semi-Disk of Acceleration i o b, there is an entire Quadrant between e S and P S, where the Acceleration is caused by the Sum of the Parallaxes. But from b to e, and from P to i, it is the Effect of their *Difference*; between b S and S e the Parallax of Longitude is positive, and prevails over the negative Effect of Latitude; but between S P and S i, the Parallax of Longitude is negative, and less than that of Latitude which is affirmative. And the same holds good with regard to the Retardation in the Semi-disk i r b, as we have in Part already observed.

4204. From the Whole of this Process it appears, that this orthographic Projection is nothing but the Praxis on the Globe, as directed for exhibiting the various Phænomena of the beginning of the Transit (4139), here projected upon a Plane; and by adapting this Projection to the *Middle* and *End* of the Transit, it will equally express, or rather give the Rationale of all the Affections of those Phases which were deduced from the Globe itself; and this I shall recommend to the Reader as a Practice that will much facilitate his Conception of these intricate Matters, and give them a lasting Impression on his Mind.

CHAP. XII.

Of the TABLES necessary for computing the TIME in which any given QUANTITY of MOTION is performed; and the MOTION produced in any given TIME; illustrated by EXAMPLES of their Use in the Doctrine of TRANSITS, and other PARTS of ASTRONOMY.

4205. **T**HERE is no Part of Astronomy which requires the Use of such Tables as serve to convert *Motion into Time*, and *Time into Motion*, more than that which is conversant about Transits, as is evident from all the preceding Calculations on this Subject. We have already explained the astronomical Principles of celestial Motions, and the chronological Doctrine of Time in the Body of these Institutes. It remains now to supply those Tables by which Motion and Time are commutable into each other, and then to explain their Uses by an Example or two.

4206. The first Table is that by which the Motion of the Earth's Equator is expressed in Time; or rather, it shews the Time in which any Number of *Degrees, Minutes, Seconds, &c.* of the Equator pass under the Meridian of a Place. This Motion is at the Rate of 15° per Hour, or one Degree in 4', because the whole 360 revolve in 24 Hours, and $24 \times 15 = 360$.

TABLE

TABLES of MOTION and TIME.

35

TABLE I.

In which the Degrees, Minutes, and Seconds of LONGITUDE, or right ASCENSION, are expressed in Hours, Minutes, and Seconds of TIME.

Deg.	H.	M.	Deg.	H.	M.	Deg.	Hours.
Min.	M.	S.	Min.	M.	S.	Deg.	
Sec.	S.	T.	Sec.	S.	T.		
1	0	4	31	2	4	70	4 40
2	0	8	32	2	8	80	5 20
3	0	12	33	2	12	90	6 0
4	0	16	34	2	16	100	6 40
5	0	20	35	2	20	110	7 20
6	0	24	36	2	24	120	8 0
7	0	28	37	2	28	130	8 40
8	0	32	38	2	32	140	9 20
9	0	36	39	2	36	150	10 0
10	0	40	40	2	40	160	10 40
11	0	44	41	2	44	170	11 20
12	0	48	42	2	48	180	12 0
13	0	52	43	2	52	190	12 40
14	0	56	44	2	56	200	13 20
15	1	0	45	3	0	210	14 0
16	1	4	46	3	4	220	14 40
17	1	8	47	3	8	230	15 20
18	1	12	48	3	12	240	16 0
19	1	16	49	3	16	250	16 40
20	1	20	50	3	20	260	17 20
21	1	24	51	3	24	270	18 0
22	1	28	52	3	28	280	18 40
23	1	32	53	3	32	290	19 20
24	1	36	54	3	36	300	20 0
25	1	40	55	3	40	310	20 40
26	1	44	56	3	44	320	21 20
27	1	48	57	3	48	330	22 0
28	1	52	58	3	52	340	22 40
29	1	56	59	3	56	350	23 20
30	2	0	60	4	0	360	24 0

4207. The Use of this Table is easy by the Example following. Let it be required to find what Time is required for the Motion of $108^{\circ} 4' 45''$ under the Meridian?

Then against	100 ^a	is	h	'	"
			6	40	0
	8 ^a	—	0	33	0
	4'	—	0	0	16
	45''	—	0	0	3

The Answer is ————— 7 12 19 (See 4074.)

4208. By the second Table you find the Motion in the Equator, or Equinoctial, corresponding to any given Time, as is plain by the Example.

TABLE II.

Shewing the MOTION corresponding to any given TIME.

Hours.	Degrees.	Min.	Deg. Min.		Min.	Deg. Min.	
		Sec.	Min.	Sec.	Sec.	Min.	Sec.
		Th.	Sec.	Th.	Th.	Sec.	Th.
1	15	1	0	15	31	7	45
2	30	2	0	30	32	8	0
3	45	3	0	45	33	8	15
4	60	4	1	0	34	8	30
5	75	5	1	15	35	8	45
6	90	6	1	30	36	9	0
7	105	7	1	45	37	9	15
8	120	8	2	0	38	9	30
9	135	9	2	15	39	9	45
10	150	10	2	30	40	10	0
11	165	11	2	45	41	10	15
12	180	12	3	0	42	10	30
13	195	13	3	15	43	10	45
14	210	14	3	30	44	11	0
15	225	15	3	45	45	11	15
16	240	16	4	0	46	11	30
17	255	17	4	15	47	11	45
18	270	18	4	30	48	12	0
19	285	19	4	45	49	12	15
20	300	20	5	0	50	12	30
21	315	21	5	15	51	12	45
22	330	22	5	30	52	13	0
23	345	23	5	45	53	13	15
24	360	24	6	0	54	13	30
		25	6	15	55	13	45
		26	6	30	56	14	0
		27	6	45	57	14	15
		28	7	0	58	14	30
		29	7	15	59	14	45
		30	7	30	60	15	0

4209. Let it be required to find what Motion in the Equator is performed in 6 Hours 20 Minutes, and 38 Seconds of Time.

		o	'	"
Then againſt 6h	is	90	0	0
20'	—	5	0	0
38"	—	0	9	30

The Answer — 95 9 30 (See 4102.)

4210. In Books of Geography you find TABLES of the LATITUDE and LONGITUDE of Places. By taking the Difference of Longitude between any two Places you find the Time corresponding thereto, and conſequently, if the Time be given in one Place, you know the Time in the other. For Example, if the Longitude of TORNEA in *Lapland* be $23^{\circ} 48' 45''$ Eaſt of *London*; then by Table I. that Motion in Longitude is made in $1^h 35' 15''$ of Time; and conſequently, when it is Noon with us, it is $35' 15''$ paſt One o'Clock there.

4211. In TABLES of ASTRONOMY, you find thoſe of the right ASCENSION and DECLINATION of the Stars; and alſo of the Sun for every Day in the Year. Now the Difference of the right Aſcenſion of the Sun, and any particular Star turned into Time, ſhews the Diſtance of Time at which the Sun and Star come upon the Meridian. For Example: On the 3d Day of *May*, the Sun's right Aſcenſion is $40^{\circ} 56' = 2^h 43' 44''$, and in the preſent Age, that of the bright Star of *Aries* is $28^{\circ} 23' = 1^h 53' 13''$; the Difference in Time is $50' 31''$, and ſo much ſooner will the Star be upon the Meridian than the Sun, or before Noon on that Day. The right Aſcenſion of ALDABARAN is $65^{\circ} 30' 3'' = 4^h 21' 17''$ this Exceſs above that of the Sun, gives $2^h 6' 33''$, and by that interval of Time it will be upon the Meridian after the Sun.

C H A P. XIII.

The DESCRIPTION and Use of a TRANSIT GLOBE, by which all the PHASES of the TRANSIT may be instantly exhibited for any given TIME and PLACE, by a new Apparatus for that Purpose.

4212. **H**AVING mentioned a TRANSIT GLOBE, which I imagined would be much more serviceable than a MAP for facilitating the Ideas relative to the Nature and various *geographical Phænomena* of the Transit, and therefore as such recommended it to the curious in the Astronomy and Geography of Transits; it may, perhaps, be necessary to give here a little further Account of the Construction and Use of such a Globe; which, after what has been already said, will require but few Words.

4213. And in order that what I have to say may be better understood, I shall explain it by Fig. 6. Plate III. where P is the North Pole of the World, and the Center of the Circle G H I described about it at the Distance $PG = 22^{\circ} 27' =$ Sun's Declination. The Globe being rectified for the *beginning and end of the Transit*, and the *two Circles* drawn as directed (4146). Then part of the first Circle is here denoted by E G Q at right Angles to the Meridian G K in which the Sun then is, *viz.* at the *beginning of the Transit*; and F H O represents part of the 2d Circle at right Angles to the Meridian H L, wherein the Sun is when the *Transit ends*.

4214. But besides these two Circles there is upon the Transit Globe a third Circle, part whereof is shewn by the Arch M C N at right Angles to the Meridian C P I, which the Sun possesses at the *middle of the Transit*. These three Circles divide the Surface of the Globe into two large Gores, and four small ones; they are distinguished from each on the small *three-inch* pocket Globes by different

rent transparent Colours; but upon the larger Globe of twelve Inches Diameter, only by coloured Circles; and then by the Cast of an Eye it will be seen.

First, What Parts of the Earth are inhabited by those who view the *whole Transit from beginning to end*, viz. those which live in the Gore, E A B F marked with a *red Colour* upon the small Globes.

Secondly, The other large Gore opposite to this is not coloured at all but appears *white*, in all which (except a little Triangle at the other End) *no part of the Transit will be visible*.

Thirdly, The first small Gore on the West Side contain'd between the Circles E A and A M, shews all those Places who see not the beginning of the Transit, but to whom the Sun will be seen to rise with the Planet from the beginning at A E, to the middle of the Transit at A M.

Fourthly, In the second small Gore contiguous to the foregoing (as contained between the Circles M B and O B,) are contained all those Parts of the Earth to whom the Sun will be seen to rise with Venus upon its Disk from the *middle of the Transit* at M B, *to the end* in the Circle O B.

Fifthly, On the other (or Eastern) Side the Globe are the other two small Gores; in the farthest of which, between the Circles A N and A Q, are contained all those Parts of the World where the Sun will set with Venus upon its Disk from the *beginning to the middle*.

Sixthly, In the small Gore between the Circles B N and B F, you will see all Parts of the Earth's Surface where the Sun sets with the Planet upon its Disk, from the *middle of the Transit to the end*.

Seventhly, The small triangular Space A B D, shews all those Places who will see the *beginning and end of the Transit*, but will be gradually deprived of the other Phases from the beginning to the middle which they can see nothing at all of, the Triangle A D B C being then intirely under the Horizon M C N. After this, they recover Sight

of the Sun and the Transit by Degrees, till all observe it at the end.

Eighthly, At the farther End of the opposite Gore (at the South Pole) is such another small Triangle, and of an equal Area, the Inhabitants of which will gradually see the Transit from the beginning to the middle, which will be seen by them all; but after that, they will by Degrees loose Sight of the Sun, till the end of the Transit becomes invisible to them all.

4215. These two Triangles are likewise distinguished by proper Colours, as also each of those *six Places* where the Transit will be observed to the greatest Advantage. (See 4152, &c.) Therefore upon the TRANSIT GLOBE you will observe every Distinction ready made to your Hand, for forming the clearest and most accurate and pleasing Ideas of the Whole, and every Part of this grand and most important of all celestial Phænomena.

4216. But what renders this Globe of the most general Use, is the *horary Quadrant of Altitude*, which by taking the Altitude of a certain Semicircle, shews the Phasis of the Transit at any given Minute of the Passage to all the Inhabitants of the illumined Surface of the Earth. The Rationale of this *Apparatus* is founded in this Consideration, that the Distance of the Semicircle E Q of the *beginning*, and the Semicircle O F of the end of the Transit, (or the Angle E D O) is equal to $86^{\circ} 4'$; and therefore if that Extent be laid down in a *Line of Hours* upon the Quadrant of Altitude, including the whole Duration of the Transit, then by it we can take the Elevation of the Semicircle E Q above the Western Horizon of the illumined Disk, and solve the following general Problem, *viz.*

4217. It is required to rectify the Globe for shewing the Appearance of the Transit at 15' past ix. (by the Time at LONDON). To do this, elevate the North Pole of the Globe $22^{\circ} 27'$ above the Horizon on the North Part, then fix the *horary Quadrant* to the Meridian in the Zenith, and bring

bring it to the Western Part of the Horizon. Lastly, elevate the Semicircle E Q to the given Hour and Minute upon the Quadrant, and there let the Globe rest; and there will be now a general Exhibition of all the Parts of the Earth where the *Transit* at the given Moment can, or cannot, be seen; as also the Time of Day to each particular Place, and the Distance the Planet is advanced in the transit Line upon the Sun's Disk.

4218. All these Things will undoubtedly appear too easy to any Person acquainted with the Use of the common *terrestrial Globe* to require any further Explication here, I shall conclude with only observing, that the same Globe (of 3 Inches Diameter) may be applied to an Orrery, so as to make a *transit* TELLURIAN, by which every thing relative to the Doctrine of *Parallaxes*, and the general *Phænomena* of the TRANSIT, may be exhibited in the most natural Manner possible.

4219. In Fig. 7. is a Representation of the Transit at *Sun-set*, as it will be observed at LONDON, if the Weather be fine. A D E is the Semi-disk of the Sun, with the Center C in the Horizon. C B is the Position of the Ecliptic, making an Angle A C B of $39^{\circ} \frac{1}{2}$; and D E is the Position of the *transit Line*, in which the Place of the Planet is at V at Sun-set.

4220. That there may be no Trouble in finding the six Places where Observations on the Transit may be made to the greatest Advantage, I have here put them together in one View, *viz.*


1. The accelerated Ingress	{ Lat. 51 28 17 N. Long. 0 6 12 E.
2. The retarded Ingress	{ Lat. 43 30 33 S. Long. 162 49 15 W.
3. The retarded Egress	{ Lat. 22 58 37 N. Long. 68 34 15 E.
4. The	


4. The accelerated Egress	{ Lat.	17 16 33	S.
	{ Long.	128 44 15	W.
5. The Duration, a <i>Maximum</i>	{ Lat.	67 49 23	N.
	{ Long.	24 20 30	E.
6. The Duration, a <i>Minimum</i>	{ Lat.	38 28 33	S.
	{ Long.	155 32 30	W.

END OF THE SECOND PART.




Plate 1.







NORTH SIDE of the ORBIT of VENUS.
Eastern Hemisphere or Morning.



NORTH SIDE of the ORBIT of ♀
Western Hemisphere or Afternoon.



SOUTH SIDE of the ORBIT of ♀
Eastern Hemisphere or Morning.



SOUTH SIDE of the ORBIT of ♀
Western Hemisphere or Afternoon.

The Sun's upper Limb

Fig. 6.




Fig. 1.



Fig. 2.




Fig. 3.




Fig. 1.




Fig. 4.




Fig. 5.

